



F.E.D.I.A.F.

NUTRITIONAL GUIDELINES FOR FEEDING PET RABBITS

31 MAY 2013

TABLE OF CONTENTS

	Sections	Content	Page
I	Acknowledgements		3
II	Glossary	Definitions	4
III	Introduction		6
IV	Basic principles	Objectives and Scope	7
		Types of Ingredient	8
		Energy intake	9
V	Nutrient recommendations	Use of the Table	11
		Table with nutrient recommendations	12
		Additional notes	13
VI	Analytical methods	Non-exhaustive list of analytical methods	19
VII	Annexes	I. Safe Production	22
		II. Labelling of Rabbit Foods	23
		III. Dietary Enrichment	24
		IV. Feeding issues	25
		V. Non-exhaustive list of feed materials	26
		VI. Bibliography	40

I. ACKNOWLEDGEMENTS

These guidelines were initiated and drafted on behalf of FEDIAF by the UK Pet Food Manufacturers' Association (PFMA) and subsequently finalised within FEDIAF and with the guidance from external experts.

PFMA and FEDIAF wishes to acknowledge the contribution of the following in writing and finalizing the guidelines:

- Erik Berenpas
- Dorothée Isler
- John Lowe
- Peter Messent
- Paul Miley
- Nicole Rabehl
- Hilke Seyffarth
- Marion Turnbull

PFMA and FEDIAF would like to thank the following external experts for giving their time to review the guidelines:

- **Professor Jürgen Zentek**, Institut für Tierernährung, Freie Universität Berlin, Germany
- **Professor Carlos De Blas**, Department of Animal Production, Universidad Politecnica Madrid, Spain
- **Professor Josef Kamphues**, Institute of Animal Nutrition, University of Veterinary Medicine Hannover, Germany
- **Professor Anna Meredith**, Royal (Dick) School of Veterinary Studies, Head of Exotic Animal and Wildlife Service, Edinburgh, United Kingdom
- **Professor Esther Plantinga**, Assistant Professor Animal Nutrition at Utrecht University, The Netherlands

The glossary contains definitions of key words used in Small Animal Nutrition in this Guideline followed by the source of the definition. Whenever appropriate, definitions are adapted to pet food.

II. GLOSSARY		
Additive (feed additive)	Substances, micro-organisms or preparations, other than feed material and premixtures, which are intentionally added to pet food or water in order to perform, in particular, one or more of the functions mentioned in Article 5(3) of regulation 1831/2003	Regulation 1831/2003 on additives for use in animal nutrition (Art 2 (2) (a)) adapted to pet food
ADF	Acid Detergent Fibre	See section V
ADL	Acid Detergent Lignin	See section V
Caecum of the rabbit	The unique musculature of the colon allows the intestinal tract of the rabbit to separate fibrous material from more digestible material; the fibrous material is passed as faeces, while the more nutritious material is passed into the caecum, then encased in a mucous covering and passed as a caecotroph (see caecotrophy).	
Caecotrophy	Process by which soft faeces (caecotrophes) are excreted and systematically re-ingested from the anus, mainly in the early morning. These caecotrophes come direct from the caecum and have a high protein (essential amino acids), vitamin and mineral content.	
Complementary pet food	Pet food which has a high content of certain substances but which, by reason of its composition, is sufficient for a daily ration only if used in combination with other foods.	Regulation 767/2009 on the marketing and use of feed (Art. 3 (1) (j)) adapted to pet food
Complete pet food	Pet food which, by reason of its composition, is sufficient for a daily ration.	Regulation 767/2009 on the marketing and use of feed (Art. 3 (1) (i)) adapted to pet food
Compound feed	Pet food consisting of a mixture of at least two feed materials, whether or not containing additives, for oral feeding in the form of complete or complementary pet food.	Regulation 767/2009 on the marketing and use of feed (Art. 3 (1) (h)) adapted to pet food
Daily ration	The average total quantity of feedingstuffs, calculated on a moisture content of 12%, required daily by an animal of a given species, age category and yield, to satisfy all its needs. The above-mentioned legal definition means the average total quantity of a specific pet food that is needed daily by a pet of a given species, age category and life style or activity to satisfy all its energy and nutrient requirements.	Regulation 1831/2003 on additives for use in animal nutrition (Art 2 (2) (f)) adapted to pet food
Digestible energy (DE)	The gross energy less the gross energy of faeces resulting from the consumption of that pet food.	McDonald et al, (1995)
Dietary enrichment	Feed or nutrients provided to the animal in addition to the manufactured or prepared diet	
DM	Dry Matter	
Feed materials	Means products of vegetable or animal origin, whose principal purpose is to meet animals' nutritional needs, in their natural state, fresh or preserved, and products derived from the industrial processing thereof, and	Regulation 767/2009 on the marketing and use of feed (Art. 3 (1) (g))

	organic or inorganic substances, whether or not containing feed additives, which are intended for use in oral animal-feeding either directly as such, or after processing, or in the preparation of compound feed, or as carrier of premixtures.	
Fibre	A generic term that refers to plants carbohydrate component that cannot be broken up by intestinal enzymes of the foregut	
Forage	Non-manufactured hay, grass and wild plants	
Maintenance energy requirement (MER)	The energy required to support energy equilibrium, (where ME equals heat production), over a long period of time.	Blaxter (1989)
Metabolizable energy (ME)	The digestible energy less the energy lost in urine and combustible gases.	McDonald et al (1995)
NRC	National Research Council (USA) is a council which is organised by the US National Academy of Sciences. The NRC Sub Committee on Rabbit Nutrition has compiled the nutrient requirements of rabbits 1977.	www.national-academies.org
NSP	Non Starch Polysaccharides	
Nutritional maximum limit	The maximum level of a nutrient in a complete pet food that, based on scientific data, has not been associated with adverse effects in healthy dogs and cats. Levels exceeding the nutritional maximum may still be safe, however, no scientific data are currently known to FEDIAF.	FEDIAF 2011
Pet food	Any product produced by a Pet food manufacturer, whether processed, partially processed or unprocessed, intended to be ingested by pet animals after placing on the market.	FEDIAF 2011
RA	Recommended Allowance.	
Recommendation	See Allowance	
SUL	Save Upper Limit	
TDF	Total Dietary Fibre	

III. INTRODUCTION

FEDIAF represents the national pet food industry associations in the EU and from Bosnia-Herzegovina, Croatia, Norway, Serbia, Switzerland and Russia, and is the spokesman for approximately 650 companies across Europe.

One of FEDIAF's main objectives is to maintain the well-being of pets by the provision of safe, well balanced and nutritionally sound pet food through its member companies. Therefore FEDIAF has compiled on the basis of work prepared by its British member PFMA the present "**Nutritional Guidelines for Feeding Pet Rabbits**" which is based on the state of the art knowledge on pet rabbit nutrition, providing pet food manufacturers with nutritional recommendations to ensure the production of well balanced and nutritionally sound diet and thus pet food for adult and growing pet rabbits. The guide does not include detailed information on reproduction, but it is hoped to do so in future. It is also important to note that this is a guide to the pet rabbit's daily diet, which can be delivered in a variety of ways (e.g. different sources of fibre).

These Guidelines are aimed at helping manufacturers developing and producing balanced rabbit nutrition for a long and healthy life of the animals.

This document is reviewed regularly and updated whenever there are new relevant technological, scientific or legislative developments in pet rabbit nutrition.

IV. BASIC PRINCIPLES

OBJECTIVES AND SCOPE

The objectives of FEDIAF's **Nutritional Guidelines for Feeding Pet Rabbits** are:

- i) To contribute to the production of nutritionally balanced pet food for rabbits, while complying with relevant EU legislation on animal nutrition. To achieve this objective, the guidelines incorporate up-to-date scientific knowledge on rabbit nutrition to:
 - Provide practical nutrient recommendations for pet food manufacturers when formulating their products for adult and growing rabbits.
 - Help pet food manufacturers to assess the nutritional value of commercial rabbit foods for healthy animals.
 - Provide guidance on some unique aspects of pet rabbit nutrition such as fibre requirements for their unique digestive tract.
- ii) To be the reference document on pet rabbit nutrition in Europe for EU and local authorities, consumer organisations, professionals, and customers.
- iii) To enhance cooperation between pet food manufacturers, pet care professionals and competent authorities by providing scientifically sound information on the formulation and assessment of pet foods for rabbits.
- iv) To complement FEDIAF's Guide to Good Practice for the Manufacture of Safe Pet Foods and the FEDIAF's Code for Good Labelling Practice for Pet Food.

FEDIAF's Nutritional Guidelines provide information on several aspects of food and nutrition:

- i) Recommendations for adequate and safe nutrient levels for the design of commercial complete and complementary foods for pet rabbits;
- ii) Tools for the assessment of the nutritional value of pet rabbit foods;
- iii) Recommendations for energy intake;
- iv) Typical ingredients suitable for use in rabbit foods;
- v) Different types of fibres: sources, and their role for rabbits;
- vi) Specific topics for rabbit nutrition, including selective feeding and supplementation of foods.

FEDIAF's Nutritional Guidelines give nutrient recommendations for healthy rabbits eating typical commercial pet foods.

- The levels in this guide reflect the amounts of essential nutrients to ensure adequate and safe nutrition in healthy individuals when consumed over its lifetime.
- These guidelines relate to pet rabbit foods manufactured from typical ingredients which are commonly used in rabbit feed.
- The nutrient levels include a safety margin for individual animal variation and nutrient interactions.
- It follows from the above statements that individual pet foods can be adequate and yet outside the recommendations, based on the manufacturer's own substantiation of nutritional adequacy and or feeding recommendations for said specific commercial food.

Unlike for other pet animals, there is also a significant market for rabbit feed produced for animals that are bred and reared for commercial purposes, including as food for humans. In terms of nutrition, some basic requirements are similar for both pet and commercial rabbits, and much information about rabbit nutrition comes from studies with commercial rabbits. Some practical issues arise, in particular with product labelling and safe manufacture ensuring that animal proteins do not enter the human food chain.

Excluded from FEDIAF's Nutritional Guidelines are foods for particular nutritional purposes (such as diets fed under veterinary supervision) and some other specialised foods.

TYPES OF INGREDIENTS

The table of nutrients given in a later section defines those nutrients known to be required in a food for pet rabbits. These nutrients need to be presented to the rabbit as a palatable food, and this section describes the more common ingredients (termed feed materials by the EU). Feed materials given on pet rabbit food labels may be listed as either individual feed materials, or in categories (defined in Directive 82/475) if the product is produced for pet rabbits only. It should be noted that use of feed materials and terms from the list given in Annex V which is taken from the EU catalogue is voluntary, and choice of other feed materials or name descriptors is permitted. The Catalogue and Annex V is indicative and producers must ensure that the raw materials used are safe. Further details are given in Annex II, under 'Labelling of Pet Rabbit Foods'.

Common feed materials and additives for rabbit foods.

Rabbits need a high level of fibre in their diet, and hay is the most common feed material for

supplying this, but several other high fibre feed materials are used, with some commonly used ones given in a non-exhaustive list in Annex V. Other feed materials are added as good sources of proteins, including soybean meal. Certain feed materials are added as source of particular nutrients, such as linseed for omega 3 fatty acids. Some feed materials such as carrots are very palatable to rabbits, but caution is required with the amount fed due to the high sugar content. Finally, pet rabbits need vitamins, trace elements and minerals. Some are presented in the natural feed materials, but many are supplemented by using a premix. Calcium may also be added separately via a premix in one of its mineral forms, such as calcium carbonate or di-calcium phosphate. Trace elements and vitamins are classified as additives, whereas sources of the major minerals such as calcium carbonate and di-calcium phosphate are classified as feed materials and not additives.

ENERGY INTAKE

A good guide for the appropriate energy intake for the pet rabbit would be:

- i. Maintenance = 100kcal ME x body weight^{0.75}. (Where body weight is expressed in kg and ME is metabolisable energy)
Alternatively
Maintenance = 0.42MJ ME x body weight^{0.75}.
- ii. Growth would be a multiple of 1.9 to 2.1 from this value.
- iii. Similarly, gestation a multiple of 1.35-2 as pregnancy progresses and lactation a multiple of 3
(Tobin, G. (1996))

It may be assumed that for a pet rabbit a daily dry matter (DM) intake of 3-5% bodyweight would be typical, however a very wide variation between individuals and according to the energy density of the diet is observed and in some cases intakes may be in excess of these values.

Prediction equations for the digestible energy (DE) content of diets tend to overestimate those with high levels of digestible fibre whilst underestimating those with added fat. Fibre plays an important role in the energy supply of the rabbit and consequently it is desirable to consider the contribution to dietary energy content in energy prediction equations. However, it is recognised that where analytical facilities are limited it may be more appropriate to use the following equation which provides reasonable practical predictions of DE.

$$\text{DE} = -1801 + 7.10\text{CP} + 12.01\text{EE} + 5.59\text{NFE}$$

Where DE is Kcal.kg⁻¹; CP is crude protein; EE is ether extract; NFE is nitrogen free extract (= 1000 - CP - EE - Crude Fibre - Ash - Moisture) each in g.kg⁻¹ as fed.

To convert the DE value to MJ.kg⁻¹ a multiplication of 0.004184 is required. The proportion of ME of DE for the rabbit could be taken as between 0.9 to 0.95.

Based on the following equation: ME/DE = 0.995 – 0.0048 x digestible protein g.kg⁻¹ / DE MJ.kg⁻¹

(Xiccato, G. And Trocino, A. (2010) Energy and protein metabolism and requirements. In: Nutrition of the Rabbit 2nd Edition eds: C de Blas and J Wiseman).

DE remains a common and practical basis for energy calculations for rabbit feeding.

Should information or analytical facilities be available as to the fibre content of a feed or a feeding stuff then a more appropriate equation, which takes into account a proportion of the dietary fibre, may be used (Villamide et al, 2009):

$$\text{DE} = 0.013\text{CP} + 0.036\text{EE} + 0.017\text{NFC} + 0.006\text{NDF}$$

Where DE is MJ.kg⁻¹ from analysis in g.kg⁻¹; CP is crude protein, EE is ether extract, NFC = organic matter less crude protein, ether extract and NDF, NDF is neutral detergent fibre, assayed with a heat stable amylase and expressed exclusive of residual ash. Mertens, D.R. (2002) Gravimetric determination of amylase –treated NDF in feeds with refluxing in beakers or crucibles: collaborative study J AOAC 85 1217-1240; Mertens, D.R. (2003) Challenges in measuring insoluble dietary fibre. J Anim Sci 81 3233-3249; EGRAN (2001) technical note: attempts to harmonise chemical analysis of feeds and faeces for rabbit feed evaluation. World Rabbit Science 9 57-64

For a more comprehensive review of this subject a number of chapters in 'Nutrition of the Rabbit' 2nd Edition by De Blas and Wiseman (2010) could be consulted.

INRA (2004) published values for nutrient content of feeds also provide a useful "text-book" source for rabbit DE values for feed ingredients

Obesity

As with cats and dogs, obesity is an increasing issue for rabbits. This is why it is important to monitor how much food they are given and to watch both their weight and body condition. This can be done using guides such as the Rabbit Size-O-Meter produced by the UK Pet Food Manufacturers' Association (downloadable from www.pfma.org.uk). The Size-O-Meter gives a series of simple steps helping with a hands-on assessment of the rabbits shape to gauge whether it is a healthy condition or not.

USE OF THE TABLE:

The table has been compiled following a review and interpretation by practising nutritionists and veterinarians of the published literature which is, in the main, for either commercial meat and/or laboratory rabbits.

The values are suggested safe and practical amounts for the design of a complete rabbit food.

The column entitled SUL in some cases consist of levels that should not be exceeded and are indicated by an asterisk, for example Calcium and vitamin D, whereas the others are levels at which no adverse effect has been observed in rabbits, higher levels may still be safe, but no scientific data are available.

Any compound feed for rabbits should clearly state how it is to be fed. Consequently, a complementary compound feed that is to be fed with forage sources, may in itself, not comply with the values in the table. However, when nutrient intake is calculated from the feeding guide, based on the amount of compound and forage and/or additional feedstuffs to be fed in the daily ration, the complete diet should fall within these values. Indeed, for enrichment reasons, as well as adequate dental wear, appropriate amounts of additional forage should be made available at all times. Similarly, free access to drinking water should be emphasised on the pack.

V. NUTRIENT RECOMMENDATIONS

N.B. It is important to note that these nutrient recommendations are for the pet rabbit's total diet. The ration may comprise contributions from various sources including manufactured food, forage, fresh vegetables and treats.

Nutrient	UNIT	Adult Maintenance recommendations		Growth, minimum requirements *		SUL	Legal maxima
		Ranges		Ranges			
Protein	g	120 to ≤170		150	180		
Arginine	g	8-9		7	n/d		
Glycine	g	n/d		n/d	n/d		
Histidine	g	n/d		3	n/d		
Isoleucine	g	n/d		6	n/d		
Leucine	g	n/d		11	n/d		
Lysine	g	5 – 8		5**	9		
Methionine + cystine	g	5.4 – 6.5		5.5	n/d		
Phenylalanine + tyrosine	g	n/d		n/d	n/d		
Threonine	g	5.8 – 6.5		6	n/d		
Tryptophan	g	n/d		2	n/d		
Valine	g	n/d		7	n/d		
Carbohydrates		-		-	-		
Starch *	g	≤ 200		n/d	135**	See * below	
ADF	g	170		n/d	n/d		
NDF	g	300 to 450		n/d	n/d		
ADL	g	55 to n/d		n/d	n/d		
((NDF+pectins)-ADF):ADF		n/d to ≤1.3		n/d	n/d		
ADF-ADL	g	110 to n/d		n/d	n/d		
Crude Fibre *	g	140 to 250		140** to 160			
Fat	g	25 to 50		30 to 50			
Minerals	-	-		-			-
Calcium	g	5		5		10	
Phosphorus	g	4		4		9	De Blas & Wiseman 2010 Nutrition of the rabbit
Ca / P ratio		1:5 to 2:1		1:5 to 2:1			
Potassium	g	6		2 to 6		16	
Sodium	g	2		1 to 2		8	
Chloride	g	1.7		1 to 5		4.8	
Magnesium	g	0.3 to 3		0.4 to 0.7		3.5	Plamenac et al 2008, Bio Trace element Res 124 110-117
Trace elements	-	-		-			-
Copper	mg	5 to 20		3 to 6		25	25
Iodine	mg	0.4 to 0.5		n/d		2	10
Iron	mg	30 to 400		100 n/d			1250
Cobalt *	mg	0.25 mg should be supplied when vitamin B12 is limited		0.1 to 1			10
Manganese	mg	8 to 15		20 to 40		75	150
Selenium	mg	0.05 to 0.32		0.1 to n/d		.35	0.5
Zinc	mg	50 to 150		40 to n/d			150
Vitamins	-	-		-			-
Vitamin A	IU	10000 to 12000		6000 to 10000**			
Vitamin E	mg	50 to 160		50** to n/d			
Vitamin D	IU	800 to 1000		500 to n/d		2000	2000
Vitamin C	mg	n/d	400	n/d		2000	Potentially a prooxidant if vitamin E levels are not comparable to vitamin C addition (Chen 1989 In vivo 3 199-209)
Vitamin K	mg	1	2	n/d			
B-group *	mg	n/d	n/d	n/d	n/d		

* please refer to Additional Notes on these issues in particular

The values in the table:

- Concerning starch; whilst it is recognised that an upper starch limit may be unnecessary providing that adequate fibre fractions are supplied, the purpose of these recommendations is to provide a safe and efficacious guide to complete diet. An upper starch provides additional safety to the diet for a pet rabbit where, despite appropriate feeding guides, owners may inadvertently overfeed the compound fraction of the daily ration and thus inadvertently provide an imbalance of starch and fibre.
- Further, upper values for fibre fractions are provided as it is recognised that excessive inappropriate fibre fraction intake could reduce the energy supply thus compromise the health and well-being of the rabbit

The values in the ranges are recommended values for typical diets based on the current best knowledge derived from published information including several scientific papers for adult rabbits, and the reference given below for growing rabbits. The limited published information accounts for some apparent discrepancies between the adult and growth values. In the future, some recommended values are likely to change as more information is published.

Growth:

These data come from limited research and we would encourage further work in this area. The majority of the data is taken from Kamphues et al (2009). Additional data comes from De Blas and Wiseman (2010) and Lebas (2000) (see comment marked “***”).

Legal maxima:

These list those additives with a maximum legal limit as authorised by the European Union. Maximum permitted levels have been determined by the legislator for several nutrients if added as a nutritional additive (i.e. trace-elements & vitamin D) (legal maximum). They are laid down in the Community Register of Feed Additives pursuant to Regulation 1831/2003/EC of the Parliament and the Council, concerning additives in feedstuffs. The legal maximum levels apply to all life stages (EU Regulation 1831/2003 in conjunction with EU register of feed additives). A legal maximum only applies when the particular trace-element or vitamin is added to the recipe as an additive, but relates to the ‘total’ amount present in the finished product [amount coming from the additive + amount from feed materials (ingredients)]. If the nutrient comes exclusively from feed materials, the legal maximum does not apply, instead the nutritional maximum, when included in the relevant tables, should be taken into account.

Fibre:

For pet rabbits, whilst it is entirely plausible to manufacture complete feeds with appropriate amounts of dietary carbohydrate (“fibre” and “starch”) to meet the nutritional and gastrointestinal physical and physiological needs, the physical nature of the fibre present in the product is the most important factor for dental wear rather than its chemical nature. An appropriate statement as to the need to supply this form of material should appear on all rabbit feed packs along the lines of “For the health and well-being of your rabbit good quality hay should be fed at all times”.

Quantification of fibre:

Fibre is essential to the rabbit on a daily basis, to maintain normal gut function, dental wear, and to provide substrate for normal fermentation in the caecum; all these factors are key to nutrient supply, normal health and behaviour in the rabbit.

The term fibre embraces a wide range of plant components and it is important key to quantify the minimum amounts and relative proportions of these that are important to the rabbit.

Fibre comes, in the main, from plant cell walls. Plant cell walls are not uniform and are extremely complex structures. Their type, size and shape depend upon the functionality of that cell wall within the plant. This in turn affects the functionality of the material in the rabbit and how it will be metabolized in the rabbit’s gut.

In general, plant cell walls are a series of polysaccharides often associated with glycoproteins, phenolic compounds, acetic acid and in some cells lignin.

A young growing plant cell has a primary cell wall, which contains few cellulosic fibrils and some non-cellulosic components. During ageing, the plant develops a secondary cell wall consisting of mainly cellulose and lignin.

In simple terms, primary cell wall (fermentable, sometimes referred to as digestible, fibre) supports the caecal fermentation needs, whilst secondary cell wall (often termed indigestible fibre) maintains gut motility and function. In absence of structured indigestible fibre, diarrhoea may develop. Plant cell walls (lignin, cellulose and silicate phytoliths) are also important in enhancing salivation and in providing dental abrasion and maintaining normal dental occlusion through extended chewing time, appropriate chewing patterns and force of the chew action.

Fibre measurement is complex and remains incomplete. Further, the terminology is also

often confusing and inconsistent. However, an approach to the fibre needs of rabbits can broadly be based on the following categories:

Total Dietary Fibre (TDF):

Refers to the total cell wall components in a plant. TDF represents all Non-Starch Polysaccharides (NSP), pectic substances, hemicellulose, cellulose and lignin. TDF is the sum of soluble and insoluble fibres (SDF and IDF) that refer to the dietary fibre components that are soluble and insoluble in water, respectively.

Neutral Detergent Fibre (NDF):

Equates to most of the hemicellulose and lignocellulose and all the cellulose.

Acid Detergent Fibre (ADF):

Equates to cellulose and most of the lignin. Consequently NDF minus ADF represents the hemicellulose content of the diet. (AOAC, 2000, method 973.18)

Acid Detergent Lignin (ADL):

Is nearly all lignins.


Other appropriate methods for fibre fractions, currently used in commercial laboratories would be the Fibertec[®] Tectator, Rellingen, Germany

Crude Fibre (CF):

CF represents 20-90% of the lignin and 30-100% cellulose depending upon the plant ingredient being analysed. Consequently, it could be considered as a poor analytical measure on which to base sound formulation of the complete diet for a rabbit. Meredith A and Prebble J (2012) Dispelling the myths, Fibre. Vet Times August 13th 6-7

Source : Diez M. PhD thesis : Contribution to the study of dietary fibre in dog's diet : effect on faeces characteristics, nutrient digestibility and blood metabolites. Department of Nutrition, Veterinary school, University of Liège, Belgium, 1998. Pp. 26-27.

The non-starch polysaccharide composition of plant material



Note: The table shows the relative proportions of each of the chemical constituents of plants measured by each method; crude fibre, acid detergent fibre (ADF), acid detergent lignin (ADL), neutral detergent fibre (NDF) and total dietary fibre (TDF).

Consequently an estimate of pectins can be derived from $TDF - NDF$

(Hall et al, 1997)

Fibre requirements for pet rabbits:

The rabbit needs a minimum fermentable (digestible) and a minimum non-fermentable (indigestible) fibre component in the complete diet (which may include both a compound food and forage material such as hay). Rather than formulating to crude fibre, it is strongly suggested that the formulation should be based upon values for the various non-starch

polysaccharide (fibre) fractions of the diet. In practical terms values for minima / maxima for NDF¹, ADF² and lignin are recommended at the stated DE content of the diet, so as to better reflect the important balance of digestible and indigestible fibre, irrespective of its origin in terms of ingredients in the complete diet. Formulating using appropriate values for these criteria are likely to result in a crude fibre content of the complete diet of 140g/kg when the diet is enriched with other sources of fibre (such as hay) or in the region of ≥ 160 g/kg. The corollary of this is that if the NDF, ADF and lignin values are ignored or unavailable to the formulator, a reasonable “safe” value for crude fibre would be in the order of 140g/kg to 160g/kg, irrespective of the ingredient matrix of the complete diet.

Research has indicated that a minimum NDF of 300g/kg total diet, a minimum ADF of 170g/kg total diet would meet the needs of the rabbit at adult maintenance. For growing rabbits an additional constraint of a maximum starch of 135g/kg total diet is recommended, whilst for the adult 200g/kg may be acceptable or even 250g/kg providing the minimum “fibre” values are exceeded (Gidenne and Garcia, 2006).

These fibre components may come from a wide range of ingredients that are incorporated in to the feed either in the form of ground material, as in pelleted or extruded feeds, or as additional loose fibre components. In respect of ground material, evidence indicates that if the indigestible fibre fraction was finely ground (less than 1mm) then the diet behaved as if containing insufficient dietary fibre, inducing adverse caecal patterns and scouring (Pairet et al, 1986 and Bouyssou et al, 1988).

The values for the carbohydrate fractions of the diet should fall within those suggested in the table of these guidelines and should be used in conjunction with the ratio indicated. Such an approach should meet the needs of the rabbit and reduce as far as possible the likelihood of digestive upsets.

A further improvement on quantifying fibre fraction involves the addition of pectin. This follows studies indicating differing fermentation rates of diets with similar NDF values. The effect is considered to be the pectin and other soluble fibre fractions. (Hall *et al*, 1997, and Gidenne and Garcia, 2006)

Minerals:

The rabbit is particularly sensitive to calcium intake and whilst amounts as high as 19-25g/kg total diet have been fed without apparent problems, for the pet rabbit a safer upper limit

¹ Fermentation needs

² Indigestible, motility needs

would be 10g/kg.

The cobalt requirement of the rabbit is principally to ensure supply to the hind gut microflora for the synthesis of Vitamin B12. It is however recommended to include sufficient vitamin B12 into the rabbits diet, and thus to remove the need for any additional cobalt, above that supplied from background levels found in the raw materials. The intestinal microflora of the rabbit has been shown to be very efficient at utilising any cobalt present in the diet.

If however B12 is not provided, a cobalt content of 0.25 mg/kg is recommended

Vitamins:

Vitamin C has only been found to be of benefit during periods of stress, otherwise it is assumed that the rabbit is capable of synthesis from glucose. Vitamin K is only suggested as required in the absence of caecotrophy.

The B-group of vitamins is generally regarded as being supplied from synthesis within the hind gut of the rabbit. However in many cases dietary supplements or dietary fortification/enrichment of these vitamins are common place to ensure sufficient vitamin supply in case of insufficient synthesis.

VI. ANALYTICAL METHODS

In order to obtain representative results, samples have to be collected and treated according to the general principles laid down in Commission Regulation 152/2009/EC of 27 January 2009 laying down Community methods of sampling for the official control of feed, as amended, and establishing Community methods of analysis for the official control of feed, as amended (see Table 4).

The analysis of only one sample may not reflect the level declared in the average analysis of the product. To obtain a representative analysis, multiple samples coming from different batches have to be analysed. A composite sample made from multiple samples is also valid. To evaluate the results of a single-sample analysis, minimum tolerances for deviation from the declared values of analytical constituents, as foreseen in ANNEX IV of Council Regulation 767/2009/EC of 13 July 2009 on the placing on the market and use of feed should be permitted, as well as tolerances for analytical latitudes.

Table 2: Non-exhaustive list of analytical methods

NUTRIENT	METHOD REFERENCE(S)
Sampling	EU method O.J. 1981 L 246 p.32 ISO/DIS 6491
Moisture	EU method O.J. 1971 L 279 p.7 ISO /DIS 6496
Protein (crude)	EU method O.J. 1993 L 179 p.8
Arginine	EU method O.J. 1998 L 257 p.14
Histidine	EU method O.J. 1998 L 257 p.14
Isoleucine	EU method O.J. 1998 L 257 p.14
Lysine	EU method O.J. 1998 L 257 p.14
Methionine	EU method O.J. 1998 L 257 p.14
Cystine/Cysteine	EU method O.J. 1998 L 257 p.14
Phenylalanine	EU method O.J. 1998 L 257 p.14
Tyrosine	EU method O.J. 1998 L 257 p.14
Threonine	EU method O.J. 1998 L 257 p.14
Valine	EU method O.J. 1998 L 257 p.14
Tryptophan	EU method O.J. 2000 L 174 p.32 2 nd ISO/CD 13904
Fat (crude)	EU method O.J. 1998 L 257 p.14
Linoleic Acid	VDLUFA method 5.6.2 B.S.I method BS684: section 2.34 : ISO 5509-1997 AOAC 15 th ed. (1990) 969.33 & 963.22
Arachidonic Acid	VDLUFA method 5.6.2

	B.S.I method BS684: section 2.34 : ISO 5509-1997 AOAC 15 th ed. (1990) 969.33 & 963.22
Fibre (crude)	EU method O.J. 1993 L 344 p.35
Ash (crude)	EU method O.J. 1971 L 155 p.13
Calcium	EU method O.J. 1971 L 155 p.13 ISO/DIS 6869
Phosphorus	EU method O.J. 1971 L 279 p.7 ISO/DIS 6491
Potassium	EU method O.J. 1971 L 155 p.13 ISO/DIS 6869
Sodium	EU method O.J. 1971 L 155 p.13 ISO/DIS 6869
Chloride	EU method O.J. 1971 L 155 p.7 §35 LMBG L06.00-5 AOAC 14th ed. (1984) 3.069-3.070 AOAC 15th ed. (1990) 920.155 & 928.04 AOAC 16th ed. (1998) potentiometric method 50.1.10
Magnesium	EU method O.J. 1973 L 83 p.21 ISO /DIS 6869
Iron	EU method O.J. 1978 L 206 p.43 ISO/DIS 6869
Copper	EU method O.J. 1978 L 206 p.43 ISO/DIS 6869
Manganese	EU method O.J. 1978 L 206 p.43 ISO/DIS 6869
Zinc	EU method O.J. 1998 L 257 p.13 ISO/DIS 6869
Iodine	Ministry of Agriculture, Fisheries and Food (1997). Dietary intake of iodine and fatty acids. Food Surveillance Information Sheet, 127. MAFF
Selenium	The Analyst 1979, 104, 784 VDLUFA, BD III method 11.6 (1993) AOAC 16 th ed. (1998) 9.1.01
Vitamin A	EU method O.J. 2000 L 174 p.32 VDLUFA method 13.1.2 2 nd ISO/CD 14565
Vitamin D *	VDLUFA method 13.8.1 D3 AOAC 15 th ed. (1990) 982.29 BS EN 12821 : 2000
Vitamin E	EU method O.J. 2000 L 174 p.32 2 nd ISO/CD 6867 VDLUFA method 13.5.4
Vitamin K	Analytical Proceedings, June 1993, Vol. 30, 266-267 (Vit. K3)

	J. of Chrom. 472 (1989) 371-379 (Vit. K1) BS EN 14148: 2003 (Vit. K1)
Thiamine	AOAC Int. 76 , (1993) 1156-1160 and 1276-1280 AOAC Int. 77 (1994) 681-686 The Analyst, 2000, No. 125, pp 353-360 EN 14122 (2003)
Riboflavin	AOAC Int. 76 (1993) 1156-1160 and 1276-1280 AOAC Int. 77 (1994) 681-686 AOAC 16 th ed. (1998) M 940.33 The Analyst, 2000, No. 125, pp 353-360 EN 14152 (2003)
Pantothenic Acid	AOAC 945.74 /42.2.05 (1990) USP XXIII, 1995, M 91
Niacin	AOAC 944.13 /45.2.04 (1990) USP XXIII, 1995, M 441
Vitamin B6 (Pyridoxine)	AOAC 16th ed. (1998) M 985.32 EN 14663: 2005
Folic Acid	AOAC 16th ed. (1998) M 944.12 Biacore AB: Folic Acid Handbook; BR 1005-19
Biotin	USP XXI, 1986, M 88 Biacore AB: Biotin Kit Handbook; BR 1005-18
Vitamin B12	USP XXIII, 1995, M171 AOAC 952.20 Biacore AB: Vitamin B12 Handbook; BR 1004-15
Choline	AOAC Int. Vol 82, No. 5 1999 pp 1156-1162 EG-Draft 15.706/1/VI/68-D/bn
Taurine	AOAC Int. Vol. 82 No. 4, 2000 pp 784-788
Total dietary fibre (TDF)	AOAC Official Method 985.29 or 45.4.07 for Total Dietary Fibre in Food and Food Products
Insoluble fibre (IF)	AOAC Method 991.42 or 32.1.16 for the Insoluble Dietary Fibre in Food and Food Products
Soluble fibre (SF)	AOAC Official Method 993.19 or 45.4.08 for Soluble Dietary Fibre in Food and Food Products
Acid detergent fibre (ADF)	AOAC 937.18
Neutral detergent fibre (NDF)	AOAC 2002.04

Vitamin D analysis of pet foods containing levels which are approaching the minimum recommendation, say between 500 and 1000 IU/kg DM is difficult and unreliable. The detection limit for HPLC methods is approximately 3000 to 5000 IU/kg. Analysis is not required if supplementation is practised and it is unlikely that un-supplemented products with adequate levels of vitamins A and E will be deficient in vitamin D.

ANNEX I – SAFE PRODUCTION OF PET RABBIT FOOD

These Guidelines are written for pet rabbits. However, manufacturers should be aware that unlike most other domestic pets, rabbits are also reared as food animals.

Pet rabbit feed producers and industry in general considers that farmed rabbit and pet rabbit feeding follow two entire different concepts: Pet rabbits are fed for a long and healthy life and not for rapid growth for meat production.

The safe production of pet food for rabbits follows the same requirements as for all pet foods. These requirements are discussed fully in the separate **FEDIAF Guide to Good Practice for the Manufacture of Safe Pet Foods**.

ANNEX II – LABELLING OF PET RABBIT FOODS

The labelling of rabbit feed must take place according to the rules of EU Regulation 767/2009 on the Marketing and Use of Feed:

Feeding instructions

The on-pack feeding guidelines for pet rabbit food are crucial. Owners should be advised that the daily diet should include forage (at least 0.5 of the daily feed dry matter intake), suitable fresh vegetables and fresh water in addition to prepared feed. Over consumption can lead to obesity and guidance to owners (via a variety of media including websites, help lines etc.) can indicate the health implications for the animal.

Composition

EU Regulation 767/2009 has different requirement on the labelling of the composition depending if the target animal is a food-producing animal or a pet animal:

Food-producing animals: Feed materials have to be listed by their specific name in descending order by weight.

Pet animals: The composition can either be indicated by listing the feed materials in descending order by weight, or by using categories of feed materials laid down in directive 82/475.

For detailed information on all other aspects of product labelling, see the **FEDIAF Code of Good Labelling Practice for Pet Food**.

ANNEX III – DIETARY ENRICHMENT

Rabbits lacking dietary enrichment may exhibit undesirable activities (Morton et al, 1993) consistent with stereotypic behaviour (Odberg 1978). This is regarded as an indicator of poor welfare, boredom, deprivation and frustration (Mason 2006). Repetitive oral activities may represent frustrated foraging in rabbits with limited access to high fibre forages. Barren or poorly enriched environments can also lead to other undesirable behaviours (Gunn 1994, Jackson 1991, Stauffacher 1992 and Wallace 1990). Allowing activities such as chewing and foraging has been shown to have a positive effect on behaviour (Berthelsen 1999), with the provision of hay proving to be the most effective in reducing abnormal behaviour (Lidfors 1997). Providing a block of wood for gnawing can also be effective, provided it is a soft wood such as willow or horse chestnut.

A rabbit should be provided with ample opportunity to chew and forage, through the provision of quantities of high fibre forage (hay, dried grasses) and supplementary snacks or treats high in indigestible fibre (gnaw sticks or similar). In addition it is important for welfare to design a diet that encourages increased time chewing in order to increase dental wear.

Dietary enrichment is thus important for a healthy rabbit.

ANNEX IV – FEEDING ISSUES

Selective Feeding

Rabbits as foraging animals have highly developed senses of taste and smell to assist in selecting the correct dietary components in their wild environment. When kept as pets, rabbits may exhibit selective feeding of mixed diets (Harcourt-Brown, 1996) (i.e. eating certain components and avoiding others), potentially leading to ingestion of a diet low in essential nutrients. Pet rabbits offered mixed diets were shown to favour components high in starch and low in calcium and therefore may be at risk of poor bone and tooth quality if permitted to feed this way (Harcourt-Brown, 1996). In addition, the high starch components are lower in fibre such that uncontrolled selective feeding or a diet lacking supplementary fibre, such as hay, can lead to poor tooth quality, insufficient dental wear and digestive conditions.

Pet owners may not be aware of the occurrence of selective feeding as food components may be scattered around the animal's environment, mixed with bedding. Where companies make mixed diets, they can optimise such products by ensuring the recipe is high in fibre ingredients, has a suitable calcium content, is well balanced for palatability of components and that clear instructions on correct feeding (not feeding more until the previous portion is finished) are on pack, so as to reduce or eliminate selective feeding (Cheeke 1994).

ANNEX V – FEED MATERIALS

**Excerpts from COMMISSION REGULATION (EU) No 68/2013 creating the Catalogue of feed materials
(only those feed materials of relevance for rabbits were taken from the Catalogue; the list is indicative and
manufacturers shall apply due diligence as to the safety of the feed materials)**

Non-exclusive list of the main feed materials

1. Cereal grains and products derived thereof

Number	Name	Description
1.1.1	Barley	Grains of <i>Hordeum vulgare</i> L.
1.1.2	Barley, puffed	Product obtained from milled or broken barley by means of a treatment in humid, warm conditions and under pressure.
1.1.3	Barley, roasted	Product of barley roasting process which is partially roasted with low colour.
1.1.4	Barley flakes	Product obtained by steaming or infra red micronizing and rolling dehusked barley. It may contain a small proportion of barley husks.
1.1.5	Barley fibre	Product of barley starch manufacture. It consists of particles of endosperm and principally of fibre.
1.1.6	Barley hulls	Product of ethanol-starch manufacture after dry milling, screening and dehulling of barley grains.
1.1.7	Barley middlings	Product obtained during the processing of screened, dehusked barley into pearl barley, semolina or flour. It consists principally of particles of endosperm with fine fragments of the outer skins and some grain screenings.
1.1.8	Barley protein	Product from barley obtained after starch and bran separation. It consists principally of protein.
1.1.9	Barley protein feed	Product from barley obtained after starch separation. It consists principally of protein and particles of endosperm.
1.1.10	Barley solubles	Product from barley obtained after wet protein and starch extraction.
1.1.11	Barley bran	Product of flour manufacture, obtained from screened grains of dehusked barley. It consists principally of fragments of the outer skins and of particles of grain from which the greater part of the endosperm has been removed.
1.1.12	Liquid barley starch	Secondary starch fraction from the production of starch from barley.
1.1.13	Malting barley screenings	Product from mechanical screening (size fractionation) consisting of undersized barley kernels and fractions of barley kernels separated before the malting process.
1.1.14	Malting barley and malt fines	Product consisting of fractions of barley kernels and malt separated during the production of malt.
1.1.15	Malting barley husks	Product from malting barley cleaning consisting of fractions of husk and fines.
1.1.16	Barley distillers solids, wet	Product of ethanol manufacture from barley. It contains solid feed fraction from distillation.
1.1.17	Barley distillers solubles, wet	Product of ethanol manufacture from barley. It contains soluble feed fraction from distillation.
1.1.18	Malt	Product from germinated cereals, dried, milled and/or extracted.
1.1.19	Malt rootlets	Product from malting cereals germination and malt cleaning consisting of rootlets, cereal fines, husks and small broken malted cereal grains. It may be milled.
1.2.1	Maize	Grains of <i>Zea mays</i> L. <i>ssp. mays</i> .
1.2.2	Maize flakes	Product obtained by steaming or infra red micronizing and rolling dehusked maize. It may contain a small proportion of maize husks.
1.2.3	Maize middlings	Product of the manufacture of flour or semolina from maize. It consists principally of fragments of the outer skins and of particles of grain from which less of the endosperm has been removed than in maize bran. It may contain some maize germ fragments.
1.2.4	Maize bran	Product of the manufacture of flour or semolina from maize. It consists principally of outer skins and some maize germ fragments, with some endosperm particles.
1.2.5	Maize cobs	Central core of a maize ear. It comprises unseparated rachis, grain and leaves.
1.2.6	Maize screenings	Fraction of maize kernels separated by the screening process at product intake.
1.2.7	Maize fibre	Product of the manufacture of maize starch. It consists principally of fibre.
1.2.8	Maize gluten	Product of the manufacture of maize starch. It consists principally of gluten obtained during separation of starch.
1.2.9	Maize gluten feed	Product obtained during the manufacture of maize starch. It is composed of bran and maize solubles. The product may also include broken maize and residues from the oil extraction of maize germs. Other products derived from starch and from the refining or fermentation of starch products may be added.
1.2.10	Maize germ	Product of the manufacture of semolina, flour or starch from maize. It consists predominately of maize germ, outer skins and parts of the endosperm.
1.2.11	Maize germ expeller	Product of oil manufacture obtained by pressing of processed maize germ to which parts of the endosperm and testa may still adhere.

1.2.12	Maize germ meal	Product of oil manufacture, obtained by extraction of processed maize germ.
1.2.13	Crude maize germ oil	Product obtained from maize germ.
1.2.14	Maize, puffed	Product obtained from milled or broken maize by means of a treatment in humid, warm conditions and under pressure.
1.2.15	Maize steep liquor	Concentrated liquid fraction from the steeping process of corn.
1.2.16	Sweet corn silage	By-product of the sweet-corn processing industry, composed of centre cobs, husks, base of the kernels, chopped and drained or pressed. Generated by chopping the sweet-corn cobs, husks and leaves, with presence of sweet-corn kernels.
1.2.17	Crushed degerminated (degermed) Maize	Product obtained by degermination of crushed maize. It consists principally of endosperm fragments and may contain some maize germ and outer skin particles.
1.3.1	Millet	Grains of <i>Panicum miliaceum</i> L.
1.4.1	Oats	Grains of <i>Avena sativa</i> L. and other cultivars of oats.
1.4.2	Dehulled oats	Dehulled grains of oats. It may be steam treated.
1.4.3	Oat flakes	Product obtained by steaming or infra red micronizing and rolling dehulled oats. It may contain a small proportion of oat husks.
1.4.4	Oat middlings	Product obtained during the processing of screened, dehulled oats into oat groats and flour. It consists principally of oat bran and some endosperm.
1.4.5	Oat bran	Product of flour manufacture, obtained from screened grains of dehulled oat. It consists principally of fragments of the outer skins and of particles of grain from which the greater part of the endosperm has been removed.
1.4.6	Oat hulls	Product obtained during dehulling of oat grains.
1.4.7	Oat, puffed	Product obtained from milled or broken oat by means of a treatment in humid, warm conditions and under pressure.
1.4.8	Oat groats	Cleaned oats with the hull removed.
1.4.9	Oat flour	Product obtained by milling of oat grains.
1.4.10	Fodder oat flour	Oats product with high content in starch, after decortication.
1.4.11	Oat feed	Product obtained during the processing of screened, dehulled oats into oat groats and flour. It consists principally of oat bran and some endosperm.
1.5.1	Quinoa seed, extracted	Cleaned whole seed of the quinoa plant (<i>Chenopodium quinoa</i> Willd.) from which the saponin contained in the seeds outer layer has been removed.
1.6.1	Broken rice	Part of rice kernel of <i>Oryza Sativa</i> L with a length less than three-quarters of a whole kernel. The rice may have been parboiled.
1.6.2	Milled rice	Husked rice from which almost all the bran and embryo have been removed during rice milling. The rice may have been parboiled.
1.6.3	Pre-gelatinized rice	Product obtained from milled or broken rice by pregelatinisation..
1.6.4	Extruded rice	Product obtained by extruding rice flour.
1.6.5	Rice flakes	Product obtained by flaking pregelatinized rice kernels or broken kernels.
1.6.6	Husked rice	Paddy (<i>Oryza Sativa</i> L.) from which the husk only has been removed. It may be parboiled. The processes of husking and handling may result in some loss of bran.
1.6.7	Ground fodder rice	Product obtained by grinding fodder rice, consisting either of green, chalky or unripe grains, sifted out during the milling of husked rice, or of normal husked grains which are yellow or spotted.
1.6.8	Rice flour	Product obtained by grinding milled rice. The rice may have been parboiled.
1.6.9	Husked rice, flour	Product obtained by grinding husked rice. The rice may have been parboiled.
1.6.10	Rice bran	Product obtained during rice milling, mainly consisting of the outer layers of the kernel (pericarp, seed coat, nucleus, aleurone) with part of the germ. The rice may have been parboiled or extruded.
1.6.11	Rice bran with calcium carbonate	Product obtained during rice milling, mainly consisting of the outer layers of the kernel (pericarp, seed coat, nucleus, aleurone) with part of the germ. It may contain up to 23% of calcium carbonate used as processing aid. The rice may have been parboiled.
1.6.12	Defatted rice bran	Rice bran resulting from oil extraction.
1.6.13	Rice bran oil	Oil extracted from stabilized rice bran.
1.6.14	Rice middlings	Product of rice flour and starch production, obtained by dry or wet milling and sieving. It consists principally of starch, protein, fat and fibre. The rice may have been parboiled. May contain up to 0.25% sodium and up to 0.25% sulphate.
1.6.15	Rice middlings with calcium carbonate	Product obtained during rice milling, mainly consisting of particles of aleurone layer and endosperm, It may contain up 23% of calcium carbonate used as processing aid. The rice may have been parboiled.
1.6.17	Rice germ	Product obtained during rice milling, mainly consisting of the embryo.
1.6.18	Rice germ expeller	Product remaining after rice germ has been crushed to expel the oil.
1.6.20	Rice protein	Product of rice starch production, obtained by wet milling sieving, separation, concentration and drying.
1.6.21	Liquid rice feed	Concentrated liquid product of wet milling and sieving rice.
1.6.22	Rice, puffed	Product obtained by expanding rice kernels or broken kernels.
1.6.23	Rice, fermented	Product obtained by fermentation of rice.
1.6.24	Malformed rice, milled /Chalky rice, milled	Product obtained during rice milling, mainly consisting of malformed kernel and/or chalky kernel and/or damaged kernel, whole or broken. It may be parboiled.
1.6.25	Immature rice, milled	Product obtained during rice milling, mainly consisting of immature and/or chalky kernel.
1.7.1	Rye	Grains of <i>Secale cereale</i> L.
1.7.2	Rye middlings	Product of flour manufacture, obtained from screened rye. It consists principally of

		particles of endosperm, with fine fragments of the outer skins and some miscellaneous parts of the grain.
1.7.3	Rye feed	Product of flour manufacture, obtained from screened rye. It consists principally of fragments of the outer skins, and of particles of grain from which less of the endosperm has been removed than in rye bran.
1.7.4	Rye bran	Product of flour manufacture, obtained from screened rye. It consists principally of fragments of the outer skins, and of particles of grain from which most of the endosperm has been removed.
1.8.1	Sorghum; [Milo]	Grains/seeds of <i>Sorghum bicolor</i> (L.) Moench
1.8.2	Sorghum white	Grains of white Sorghum
1.8.3	Sorghum gluten feed	Dried product obtained during the separation of sorghum starch. It consists principally of bran and a small quantity of gluten. The product may also include dried residues of maceration water and germs could be added.
1.9.1	Spelt	Grains of spelt <i>Triticum spelta</i> L., <i>Triticum dicoccum</i> Schrank, <i>Triticum monococcum</i> .
1.9.2	Spelt bran	Product of the manufacture of spelt flour. It consists principally of outer skins and some spelt germ fragments, with some endosperm particles.
1.9.3	Spelt hulls	Product obtained during dehulling of spelt grains.
1.9.4	Spelt middlings	Product obtained during the processing of screened, dehulled spelt into spelt flour. It consists principally of particles of endosperm with fine fragments of the outer skins and some grain screenings.
1.10.1	Triticale	Grains of <i>Triticum X Secale cereale</i> L. Hybrid.
1.11.1	Wheat	Grains of <i>Triticum aestivum</i> L., <i>Triticum durum</i> Desf. and other cultivars of wheat.
1.11.2	Wheat rootlets	Product from malting wheat germination and malt cleaning consisting of rootlets, cereal fines, husks and small broken malted wheat grains.
1.11.3	Wheat, pre-gelatinised	Product obtained from milled or broken wheat by means of a treatment in humid, warm conditions and under pressure.
1.11.4	Wheat middlings	Product of flour manufacture obtained from screened grains of wheat or dehulled spelt. It consists principally of particles of endosperm with fine fragments of the outer skins and some grain screenings.
1.11.5	Wheat flakes	Product obtained by steaming or infra red micronizing and rolling dehulled wheat. It may contain a small proportion of wheat husks.
1.11.6	Wheat feed	Product of flour or malting manufacture obtained from screened grains of wheat or dehulled spelt. It consists principally of fragments of the outer skins and of particles of grain from which less of the endosperm has been removed than in wheat bran.
1.11.7	Wheat bran	Product of flour or malting manufacture obtained from screened grains of wheat or dehulled spelt. It consists principally of fragments of the outer skins and of particles of grain from which the greater part of the endosperm has been removed.
1.11.8	Malted fermented wheat particles	Product obtained by a process combining malting and fermentation of wheat and wheat bran. The product is then dried and ground.
1.11.10	Wheat fibre	Fibre extracted from wheat processing. It consists principally of fibre.
1.11.11	Wheat germ	Product of flour milling consisting essentially of wheat germ, rolled or otherwise, to which fragments of endosperm and outer skin may still adhere.
1.11.12	Wheat germ, fermented	Product of fermentation of wheat germ, with inactivated micro-organisms.
1.11.13	Wheat germ expeller	Product of oil manufacture, obtained by pressing of wheat germ (<i>Triticum aestivum</i> L., <i>Triticum durum</i> Desf. and other cultivars of wheat and dehulled spelt (<i>Triticum spelta</i> L., <i>Triticum dicoccum</i> Schrank, <i>Triticum monococcum</i> L.)) to which parts of the endosperm and testa may still adhere.
1.11.15	Wheat protein	Wheat protein extracted during starch or ethanol production, maybe partially hydrolysed
1.11.16	Wheat gluten feed	Product of the manufacture of wheat starch and gluten. It consists of bran, from which the germ may have been partially removed. Wheat solubles, broken wheat and other products derived from starch and from the refining or fermentation of starch products may be added.
1.11.18	Vital wheat gluten	Wheat protein characterized by a high viscoelasticity as hydrated, with minimum 80% protein (Nx6.25) and maximum 2% ash on dry substance.
1.11.19	Liquid wheat starch	Product obtained from the production of starch/glucose and gluten from wheat.
1.11.20	Wheat starch containing protein, partially de-sugared	Product obtained during the production of wheat starch mainly comprising partially sugared starch, the soluble proteins and other soluble parts of the endosperm.
1.11.21	Wheat solubles	Product of wheat obtained after wet protein and starch extraction. May be hydrolysed.
1.11.22	Wheat yeast concentrate	Wet by-product that is released after the fermentation of wheat starch for alcohol production.
1.11.23	Malting wheat screenings	Product from mechanical screening (size fractionation) consisting of undersized wheat kernels and fractions of wheat kernels separated before the malting process.
1.11.24	Malting wheat and malt fines	Product consisting of fractions of wheat kernels and malt separated during the production of malt.
1.11.25	Malting wheat husks	Product from malting wheat cleaning consisting of fractions of husk and fines.
1.12.2	Grain flour	Flour from milling grains.
1.12.3	Grain protein concentrate	Concentrate and dried product obtained from grain after starch removing through yeast fermentation.
1.12.4	Cereal grains screenings	Products from mechanical screening (size fractionation) consisting of small grains and fractions of grain kernels, which may be germinated, separated before further processing of the grain. The products contain more crude fibre (e.g. hulls) than the unfractionated cereals

1.12.5	Grain germ	Product of flour milling and the manufacture of starch consisting principally of grain germ, rolled or otherwise, to which fragments of endosperm and outer skin may still adhere.
1.12.6	Grain spent wash syrup Error! Bookmark not defined.	Product of grain obtained through the evaporation of the concentrate of the spent wash from the fermentation and distillation of grain used in the production of grain spirit.
1.12.7	Moist distillers' grains	Moist product produced as the solid fraction by centrifuging and/or filtration of the spent wash from fermented and distilled grains used in the production of grain spirit.
1.12.8	Concentrated Distillers Solubles	Moist product from production of alcohol by fermentation and distilling a mash of wheat and sugar syrup after previous separation of bran and gluten. They may contain dead cells and/or parts of the fermentation microorganisms.
1.12.9	Distillers' grains and solubles	Product obtained when producing alcohol by fermentation and distilling grain mash of cereals and/ or other starchy and sugar containing products. They may contain dead cells and/or parts of the fermentation microorganisms. May contain 2% sulphate.
1.12.10	Distillers' dried grains	Product of alcohol distilling obtained by drying solid residues of fermented grains.
1.12.11	Distillers' dark grains; [Distillers' dried grains and solubles]	Product of alcohol distilling obtained by drying solid residues of fermented grains to which pot ale syrup or evaporated spent wash has been added.
1.12.12	Brewers' grains	Product of brewing composed by residues of malted and unmalted cereals and other starchy products, which may contain hop materials. Typically marketed in a moist condition but may also be sold in a dried form. May contain up to 0.3% dimethyl polysiloxane, may contain up to 1.5% enzymes, may contain up to 1.8% bentonite.
1.12.13	Draff	Solid product of cereal whisky production. It consists of the residues from hot water extraction of malted cereal. Typically marketed in the moist form after the extract has been removed by gravity.
1.12.14	Mash Filter Grains	Solid product obtained through the production of beer, malt extract and whisky spirit. It consists of the residues of hot water extraction of ground malt and possibly other sugar or starch-rich adjuncts. Typically marketed in the moist form after the extract has been removed by pressing.
1.12.15	Pot ale	The product remaining in the still from the first (wash) distillation of a malt distillery.
1.12.16	Pot ale syrup	Product from the first (wash) distillation of a malt distillery produced by evaporating the pot ale remaining in the still.

2. Oil seeds, oil fruits, and products derived thereof

Number	Name	Description
2.1.1	Babassu expeller	Product of oil manufacture, obtained by pressing Babassu palm nuts <i>Orbignya</i> varieties.
2.2.1	Camelina seed	Seeds of <i>Camelina sativa</i> L. Crantz.
2.2.2	Camelina, expeller	Product of oil manufacture, obtained by pressing of seeds of Camelina.
2.2.3	Camelina meal	Product of oil manufacture, obtained by extraction and appropriate heat treatment of Camelina seed expeller.
2.3.1	Cocoa husks	Teguments of the dried and roasted beans of <i>Theobroma cacao</i> L.
2.3.2	Cocoa hulls	Product obtained by processing of cocoa beans.
2.3.3	Cocoa bean meal, partially decorticated	Product of oil manufacture, obtained by extraction of dried and roasted cocoa beans <i>Theobroma cacao</i> L. from which part of the husks has been removed
2.4.1	Copra expeller	Product of oil manufacture, obtained by pressing the dried kernel (endosperm) and outer husk (tegument) of the seed of the coconut palm <i>Cocos nucifera</i> L.
2.4.2	Copra, hydrolysed expeller	Product of oil manufacture, obtained by pressing and enzymatic hydrolysis of the dried kernel (endosperm) and outer husk (tegument) of the seed of the coconut palm <i>Cocos nucifera</i> L.
2.4.3	Copra meal	Product of oil manufacture, obtained by extraction of the dried kernel (endosperm) and outer husk (tegument) of the seed of the coconut palm.
2.5.1	Cotton seed	Seeds of <i>Gossypium</i> spp. from which the fibres have been removed.
2.5.2	Cotton seed meal, partially decorticated	Product of oil manufacture, obtained by extraction of seeds of cotton from which the fibres and part of the husks have been removed. (Maximum crude fibre 22.5% in the dry matter).
2.5.3	Cotton seed expeller	Product of oil manufacture, obtained by pressing of seeds of cotton from which the fibres have been removed.
2.6.1	Groundnut expeller, partially decorticated	Product of oil manufacture, obtained by pressing of partially decorticated groundnuts <i>Arachis hypogaea</i> L. and other species of <i>Arachis</i> . (Maximum crude fibre content 16% in the dry matter)
2.6.2	Groundnut meal, partially decorticated	Product of oil manufacture, obtained by extraction of partially decorticated groundnut expeller. (Maximum crude fibre content 16% in the dry matter)
2.6.3	Groundnut expeller, decorticated	Product of oil manufacture, obtained by pressing of decorticated groundnuts.
2.6.4	Groundnut meal, decorticated	Product of oil manufacture, obtained by extraction of decorticated groundnut expeller.
2.7.1	Kapok expeller	Product of oil manufacture obtained by pressing of Kapok seeds (<i>Ceiba pentadra</i> L. Gaertn.).
2.8.1	Linseed	Seeds of linseed <i>Linum usitatissimum</i> L. (Minimum botanical purity 93%) as whole, flattened or ground linseed.
2.8.2	Linseed expeller	Product of oil manufacture, obtained by pressing of linseed. (Minimum botanical purity

		93%).
2.8.3	Linseed meal	Product of oil manufacture, obtained by extraction and appropriate heat treatment of linseed expeller.
2.8.4	Linseed expeller feed	Product of oil manufacture, obtained by pressing of linseed. (Minimum botanical purity 93%). May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants.
2.8.5	Linseed meal feed	Product of oil manufacture, obtained by extraction and appropriate heat treatment of linseed expeller. May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants.
2.9.2	Mustard seed meal	Product obtained by the extraction of volatile mustard oil from mustard seeds
2.10.1	Niger seed	Seeds of the niger plant <i>Guizotia abyssinica</i> (L. F.) Cass.
2.10.2	Niger seed expeller	Product of oil manufacture, obtained by pressing of seeds of the niger plant (Ash insoluble in HCl: maximum 3,4%)
2.11.1	Olive pulp	Product of oil manufacture, obtained by extraction of pressed olives <i>Olea europea</i> L. separated as far as possible from parts of the kernel.
2.11.2	Defatted olive meal feed	Product of olive oil manufacture, obtained by extraction and appropriate heat treatment of olive pulp expeller separated as far as possible from parts of the kernel. May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants.
2.11.3	Defatted olive meal	Product of olive oil manufacture, obtained by extraction and appropriate heat treatment of olive pulp expeller separated as far as possible from parts of the kernel.
2.12.1	Palm kernel expeller	Product of oil manufacture, obtained by pressing of palm kernels <i>Elaeis guineensis</i> Jacq., <i>Corozo oleifera</i> (HBK) L. H. Bailey (<i>Elaeis melanococca auct.</i>) from which as much as possible of the hard shell has been removed.
2.12.2	Palm kernel meal	Product of oil manufacture, obtained by extraction of palm kernels from which as much as possible of the hard shell has been removed.
2.13.1	Pumpkin and squash seed	Seeds of <i>Cucurbita pepo</i> L. and plants of the genus <i>Cucurbita</i> .
2.13.2	Pumpkin and squash seed, expeller	Product of oil manufacture, obtained by pressing of seeds of <i>Cucurbita pepo</i> and plants of the genus <i>Cucurbita</i> .
2.14.1	Rape seed	Seeds of rape <i>Brassica napus</i> L. ssp. <i>oleifera</i> (Metzg.) Sinsk., of Indian sarson <i>Brassica napus</i> L. var. <i>glauca</i> (Roxb.) O.E. Schulz and of rape <i>Brassica rapa</i> ssp. <i>oleifera</i> (Metzg.) Sinsk. Minimum botanical purity 94%.
2.14.2	Rape seed, expeller	Product of oil manufacture, obtained by pressing of seeds of rape.
2.14.3	Rape seed meal	Product of oil manufacture, obtained by extraction and appropriate heat treatment of rape seed expeller.
2.14.4	Rape seed, extruded	Product obtained from whole rape by means of a treatment in humid, warm conditions and under pressure increasing starch gelatinisation.
2.14.5	Rape seed protein concentrate	Product of oil manufacture, obtained by separation of protein fraction of rapeseed expeller or rapeseed.
2.14.2	Rape seed expeller feed	Product of oil manufacture, obtained by pressing of seeds of rape. May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants.
2.14.3	Rape seed meal feed	Product of oil manufacture, obtained by extraction and appropriate heat treatment of rape seed expeller. May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants.
2.15.1	Safflower seed	Seeds of the safflower <i>Carthamus tinctorius</i> L.
2.15.2	Safflower seed meal, partially decorticated	Product of oil manufacture, obtained by extraction of partially decorticated seeds of safflower.
2.15.3	Safflower hulls	Product obtained during dehulling of safflower seeds.
2.16.1	Sesame seed	Seeds of <i>Sesamum indicum</i> L.
2.17.1	Sesame seed, partially dehulled	Product of oil manufacture, obtained by removing part of the husks.
2.17.2	Sesame hulls	Product obtained during dehulling of sesame seeds.
2.17.3	Sesame seed expeller	Product of oil manufacture, obtained by pressing of seeds of the sesame plant (Ash insoluble in HCl: maximum 5%)
2.18.1	Toasted soya (beans)	Soya beans (<i>Glycine max.</i> L. Merr.) subjected to an appropriate heat treatment. (Urease activity maximum 0.4 mg N/g × min.).
2.18.2	Soya (bean) expeller	Product of oil manufacture, obtained by pressing the seed of soya
2.18.3	Soya (bean) meal	Product of oil manufacture, obtained from soya beans after extraction and appropriate heat treatment. (Urease activity maximum 0.4 mg N/g × min.).
2.18.4	Soya (bean) meal, dehulled	Product of oil manufacture, obtained from dehulled soya beans after extraction and appropriate heat treatment. (Urease activity maximum 0.5 mg N/g × min.).
2.18.5	Soya (bean) hulls	Product obtained during dehulling of soya beans.
2.18.6	Soya beans, extruded	Product obtained from soya beans by means of a treatment in humid, warm conditions and under pressure increasing starch gelatinisation.
2.18.7	Soya (bean) protein concentrate	Product obtained from dehulled, fat extracted soya beans, after fermentation or a second extraction to reduce the level of nitrogen-free extract.

2.18.8	Soya bean pulp ; [Soya bean paste]	Product obtained during extraction of soya beans for food preparation.
2.18.9	Soya bean molasses	Product obtained during the processing of soya bean.
2.18.10	By-product from soybean preparation	Products obtained when processing soybeans to obtain soybean food preparations.
2.18.11	Soya (beans)	Soya beans (<i>Glycine max.</i> L. Merr.)
2.18.12	Soybean, flakes	Product obtained by steaming or infra red micronizing and rolling dehulled soya beans. (Urease activity maximum 0.4 mg N/g × min.).
2.18.13	Soya (bean) meal feed	Product of oil manufacture, obtained from soya beans after extraction and appropriate heat treatment. (Urease activity maximum 0.4 mg N/g × min.). May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants.
2.18.14	Soya (bean) meal feed, dehulled	Product of oil manufacture, obtained from dehulled soya beans after extraction and appropriate heat treatment. (Urease activity maximum 0.5 mg N/g × min.). May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants.
2.19.1	Sunflower seed	Seeds of the sunflower <i>Helianthus annuus</i> L.
2.19.2	Sunflower seed expeller	Product of oil manufacture, obtained by pressing of seeds of the sunflower.
2.19.3	Sunflower seed meal	Product of oil manufacture, obtained by extraction and appropriate heat treatment of sunflower seed expeller.
2.19.4	Sunflower seed meal, dehulled	Product of oil manufacture, obtained by extraction and appropriate heat treatment of expeller of sunflower seeds from which part or all of the husks has been removed. Maximum crude fibre 27.5% in the dry matter
2.19.5	Sunflower seed hulls	Product obtained during dehulling of sunflower seeds.
2.19.6	Sunflower seed meal feed	Product of oil manufacture, obtained by extraction and appropriate heat treatment of sunflower seed expeller. May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants.
2.19.7	Sunflower seed meal feed, dehulled	Product of oil manufacture, obtained by extraction and appropriate heat treatment of expeller of sunflower seeds from which part or all of the husks has been removed. May contain up to 1% used bleaching earth and filter aid (e.g. diatomaceous earth, amorphous silicates and silica, phyllosilicates and cellulosic or wood fibres) and crude lecithins from integrated crushing and refining plants. Maximum crude fibre 27.5% in the dry matter.
2.20.1	Vegetable oil and fat	Oil and fat obtained from plants (excluding castor oil from the ricinus plant), it may be degummed, refined and/or hydrogenated
2.21.1	Crude lecithins	Product obtained during degumming of crude oil from oilseeds and oil fruits with water. Citric acid, phosphoric acid or sodium hydroxide may be added during degumming of the crude oil.
2.22.1	Hemp seed	Controlled hemp seed <i>Cannabis sativa</i> L. with a maximum THC content according to EU legislation
2.22.2	Hemp expeller	Product of oil manufacture obtained by pressing of hemp seed.
2.22.3	Hemp oil	Product of oil manufacture, obtained by pressing of hemp plant and seed.
2.23.1	Poppy seed	Seeds of <i>Papaver somniferum</i> L.
2.23.2	Poppy meal	Product of oil manufacture, obtained by extraction of expeller of poppy seed

3. Legume seeds and products derived thereof

Number	Name	Description
3.1.1	Beans, toasted	Seeds of <i>Phaseolus</i> spp. or <i>Vigna</i> spp. submitted to an appropriate heat treatment.
3.1.2	Bean protein concentrate	Product obtained from the separated bean fruit water, when producing starch.
3.2.1	Carob, dried	Dried fruits of the carob tree <i>Ceratonia siliqua</i> L.
3.2.3	Carob pods, dried	Product obtained by crushing the dried fruits (pods) of the carob tree and from which the locust beans have been removed.
3.2.4	Dried carob pod meal, micronised	Product obtained by micronisation of the dried fruits of the carob tree from which the locust beans have been removed.
3.2.5	Carob germ	Germ of the locust bean of the carob tree.
3.2.6	Carob germ, expeller	Product of oil manufacture, obtained by pressing of germ of carob.
3.2.7	Locust bean (seed)	Bean of the carob tree.
3.3.1	Chick peas	Seeds of <i>Cicer arietinum</i> L.
3.4.1	Ervil	Seeds of <i>Ervum ervilia</i> L.
3.5.1	Fenugreek seed	Seed of fenugreek (<i>Trigonella foenum-graecum</i>).
3.6.1	Guar meal	Product obtained after extraction of the mucilage from seeds of guar bean <i>Cyamopsis tetragonoloba</i> (L.) Taub.
3.6.2	Guar germs meal	Product of mucilage extraction from the germ of seeds of guar bean.
3.7.1	Horse beans	Seeds of <i>Vicia faba</i> L. ssp. <i>faba</i> L. and var. <i>equina</i> Pers. and var. <i>minuta</i> (Alef.) Mansf.
3.7.2	Horse bean flakes	Product obtained by steaming or infra red micronizing and rolling dehulled horse beans.
3.7.3	Film horse beans; [Faba bean hulls]	Product obtained during dehulling horse bean seeds, consisting mainly of external envelopes.

3.7.4	Horse beans, dehulled	Product obtained during dehulling horse bean seeds, consisting mainly of bean kernels from horse beans.
3.7.5	Horse bean protein	Product obtained by grinding and air fractionation of horse beans.
3.8.1	Lentils	Seeds of <i>Lens culinaris</i> a.o. Medik.
3.8.2	Lentil hulls	Product obtained during dehulling process of lentil seeds.
3.9.1	Sweet lupins	Seeds of <i>Lupinus</i> spp. low in bitter seed content.
3.9.2	Sweet lupins, dehulled	Dehulled lupin seeds.
3.9.3	Film lupins; [lupin hulls]	Product obtained during dehulling of lupin seeds, consisting mainly of external envelopes.
3.9.4	Lupin pulp	Product obtained after extraction of components of lupin.
3.9.5	Lupin middlings	Product obtained during the manufacture of lupin flour from lupin. It consists principally of particles of cotyledon, and to a lesser extent, of skins.
3.9.6	Lupin protein	Product obtained from the separated lupin fruit water when producing starch, or after grinding and air fractionation.
3.9.7	Lupin protein meal	Product of lupin processing to produce a high protein meal.
3.10.1	Mung beans	Beans of <i>Vigna radiata</i> L.
3.11.1	Peas	Seeds of <i>Pisum</i> spp.
3.11.2	Pea bran	Product obtained during the manufacture of pea meal. It is composed mainly of skins removed during the skinning and cleaning of peas.
3.11.3	Pea flakes	Product obtained by steaming or infra red micronizing and rolling dehulled seeds of peas.
3.11.4	Pea flour	Product obtained during the grinding of peas.
3.11.5	Pea hulls	Product obtained during the manufacture of pea meal from peas. It is mainly composed of skins removed during the skinning and cleaning and, to a lesser extent, of endosperm.
3.11.6	Peas, dehulled	Dehulled pea seeds.
3.11.7	Pea middlings	Product obtained during the manufacture of pea flour. It consists principally of particles of cotyledon, and to a lesser extent, of skins.
3.11.8	Pea screenings	Product from the mechanical screening consisting of fractions of pea kernels separated before further processing.
3.11.9	Pea protein	Product obtained from the separated pea fruit water when producing starch, or after grinding and air fractionation, maybe partially hydrolysed.
3.11.10	Pea pulp	Product obtained from starch and protein wet extraction from peas. It is mainly composed of internal fibre and starch.
3.11.11	Pea solubles	Product obtained from starch and protein wet extraction from peas. It is mainly composed of soluble proteins and oligosaccharides.
3.11.12	Pea fibre	Product obtained by extraction after grinding and sieving of the dehulled pea.
3.12.1	Vetches	Seeds of <i>Vicia sativa</i> L. var <i>sativa</i> and other varieties.
3.13.1	Chickling vetch	Seeds of <i>Lathyrus sativus</i> L. submitted to an appropriate heat treatment.
3.14.1	Monantha vetch	Seeds of <i>Vicia monanthos</i> Desf.

4. Tubers, roots, and products derived thereof

Number	Name	Description
4.1.1	Sugar beet	Root of <i>Beta vulgaris</i> L. ssp. <i>vulgaris</i> var. <i>altissima</i> Doell.
4.1.2	Sugar beet tops and tails	Fresh product of the manufacture of sugar consisting mainly of cleaned pieces of sugar beet with or without parts of beet leaves.
4.1.3	(Beet) sugar; [sucrose]	Sugar extracted from sugar beets using water.
4.1.4	(Sugar) beet molasses	Syrupy product obtained during the manufacture or refining of sugar from sugar beets. May contain up to 0.5% antifoaming agents. May contain up to 0.5% antiscaling agents. May contain up to 2% sulphate. May contain up to 0.25% sulphite.
4.1.5	(Sugar) beet molasses, partially-desugared and/or debetainized	Product obtained after further extraction using water of sucrose and/or betaine from sugar beet molasses. May contain up to 2% sulphate. May contain up to 0.25% sulphite.
4.1.6	Isomaltulose molasses	Non-crystallised fraction from the manufacture of isomaltulose by enzymatic conversion of sucrose from sugar beets.
4.1.7	Wet (sugar) beet pulp	Product of the manufacture of sugar consisting of slices of sugar beet that have had sugar extracted with water. Minimum moisture content: 82%. Sugar content is low and declines towards zero due to (lactic acid) fermentation.
4.1.8	Pressed (sugar) beet pulp	Product of the manufacture of sugar consisting of slices of sugar beet that have had sugar extracted with water and have been mechanically pressed. Maximum moisture content: 82%. Sugar content is low and declines towards zero due to (lactic acid) fermentation. May contain up to 1% sulphate.
4.1.9	Pressed (sugar) beet pulp, molassed	Product of the manufacture of sugar consisting of slices of sugar beet that have had sugar extracted with water, have been mechanically pressed, and with molasses added. Maximum moisture content: 82%. Sugar content declines due to (lactic acid) fermentation. May contain up to 1% sulphate.
4.1.10	Dried (sugar) beet pulp	Product of the manufacture of sugar consisting of slices of sugar beet that have had sugar extracted with water, mechanically pressed and dried. May contain up to 2% sulphate.
4.1.11	Dried (sugar) beet pulp, molassed	Product of the manufacture of sugar consisting of slices of sugar beet that have had sugar extracted with water, mechanically pressed, and dried, with molasses added. May contain up to 0.5% antifoaming agents. May contain up to 2% sulphate.

4.1.12	Sugar syrup	Product obtained by processing of sugar and/or molasses. May contain up to 0.5% sulphate. May contain up to 0.25% sulphite.
4.1.13	(Sugar) beet pieces, boiled	Product of the manufacture of edible syrup from sugar beet, which may be pressed or dried.
4.1.14	Fructo-oligosaccharides	Product obtained from sugar from sugar beet through an enzymatic process.
4.2.1	Beetroot juice	Juice from pressing of red beet (<i>Beta vulgaris convar. crassa var. conditiva</i>) with subsequent concentration and pasteurization, maintaining the typical vegetable-like taste and flavour.
4.3.1	Carrots	Root of the yellow or red carrot <i>Daucus carota</i> L.
4.3.2	Carrot peelings, steamed	Moist product from the carrot processing industry consisting of the peelings removed from the carrot root by steam treatment to which auxiliary flows of gelatinous carrot starch may be added. Maximum moisture content: 97%.
4.3.3	Carrot scrapings	Moist product which is released via mechanical separation in the processing of carrots and which mostly consists of dried carrots and carrot remnants. The product may have been subject to heat treatment. Maximum moisture content: 97%.
4.3.4	Carrot flakes	Product obtained by flaking roots of the yellow or red carrot, which are subsequently dried.
4.3.5	Carrot, dried	Root of the yellow or red carrot regardless of their presentation, which are subsequently dried.
4.3.6	Carrot feed, dried	Product constituted of internal pulp and outer skins that are dried.
4.4.1	Chicory roots	Roots of <i>Cichorium intybus</i> L.
4.4.2	Chicory tops and tails	Fresh product from chicory processing. It consists predominantly of cleaned pieces of chicory and parts of leaves.
4.4.3	Chicory seed	Seed of <i>Cichorium intybus</i> L.
4.4.4	Pressed chicory pulp	Product of the manufacture of inulin from roots of <i>Cichorium intybus</i> L. consisting of extracted and mechanically pressed slices of chicory. The (soluble) chicory carbohydrates and water have been partly removed. May contain up to 1% sulphate, and may contain up to 0.2% sulphite.
4.4.5	Dried chicory pulp	Product of the manufacture of inulin from roots of <i>Cichorium intybus</i> L. consisting of extracted and mechanically pressed slices of chicory and subsequent drying. The (soluble) chicory carbohydrates have been partly extracted. May contain up to 2% sulphate, and may contain up to 0.5% sulphite.
4.4.6	Chicory roots powder	Product obtained by chopping, drying and grinding of chicory roots. May contain up to 1% of anticaking agents.
4.4.7	Chicory molasses	Product of chicory processing, obtained during the production of inulin and oligofructose. Chicory molasses consists of organic plant material and minerals. May contain up to 0.5% antifoaming agents
4.4.8	Chicory vinasses	By-product from chicory processing obtained after the separation of inulin and oligofructose and ion exchange elution. Chicory vinasses consists of organic plant material and minerals. May contain up to 1% antifoaming agents.
4.4.9	Chicory inulin	Inulin is a fructan extracted from roots of <i>Cichorium intybus</i> L.; raw chicory inulin may contain up to 1% sulphate and may contain up to 0.5% sulphite.
4.4.10	Oligofructose syrup	Product obtained by partial hydrolysis of inulin from <i>Cichorium intybus</i> L.; raw oligofructose syrup may contain up to 1% sulphate and may contain up to 0.5% sulphite.
4.4.11	Oligofructose, dried	Product obtained by partial hydrolysis of inulin from <i>Cichorium intybus</i> L. and subsequent drying.
4.5.1	Garlic, dried	White to yellow powder of pure, ground garlic, <i>Allium sativum</i> L.
4.6.1	Manioc; [tapioca]; [cassava]	Roots of <i>Manihot esculenta</i> Crantz, regardless of their presentation.
4.6.2	Manioc, dried	Roots of Manioc, regardless of their presentation, which are subsequently dried.
4.7.1	Onion pulp	Moist product which is released during the processing of onions (genus <i>Allium</i>) and consists of both skins and whole onions. If from the production process for onion oil, then it mostly consists of cooked remains of onions.
4.7.2	Onions, fried	Skinned and crumbed onion pieces which are then fried.
4.7.3	Onions solubles	Dry product which is released during the processing of fresh onions. It is obtained by alcoholic and/or water extraction, the water or alcoholic fraction is separated and spraydried. It consists mainly in carbohydrates.
4.8.1	Potatoes	Tubers of <i>Solanum tuberosum</i> L.
4.8.2	Potatoes, peeled	Potatoes from which the skin is removed using steam treatment.
4.8.3	Potato peelings, steamed	Moist product from the potato processing industry consisting of the peelings removed by steam treatment from the potato tuber to which auxiliary flows of gelatinous potato starch may be added. It may be mashed.
4.8.4	Potato cuttings, raw	Product released from potatoes during the preparation of potato products for human consumption, which may have been peeled.
4.8.5	Potato scrapings	Product which is released via mechanical separation in the processing of potatoes and which mostly consists of dried potatoes and potato remnants. The product may have been subject to heat treatment.
4.8.6	Potato, mashed	Blanched or boiled and then mashed potato product.
4.8.7	Potato flakes	Product obtained by rotary drying of washed, peeled or unpeeled steamed potatoes.
4.8.8	Potato pulp	Product of the manufacture of potato starch consisting of extracted ground potatoes.
4.8.9	Potato pulp, dried	Dried product of the manufacture of potato starch consisting of extracted ground potatoes.
4.8.10	Potato protein	Product of starch manufacture composed mainly of protein substances obtained after the

		separation of starch.
4.8.11	Potato protein, hydrolysed	Protein obtained by a controlled enzymatic hydrolysis of potato proteins.
4.8.12	Potato protein, fermented	Product obtained by fermentation of potato protein and subsequent spray drying.
4.8.13	Potato protein fermented, liquid	Liquid product obtained by fermentation of potato protein.
4.8.14	Potato juice, concentrated	Concentrated product of the manufacture of potato starch, consisting of the remaining substance after the partial removal of fibre, proteins and starch from the whole potato pulp and evaporation of part of the water.
4.8.15	Potato granules	Dried potatoes (potatoes after washing, peeling, size reduction – cutting, flaking, etc. and water content removal).
4.9.1	Sweet potato	Tubers of <i>Ipomoea batatas</i> L. regardless of their presentation.
4.10.1	Jerusalem artichoke; [Topinambur]	Tubers of <i>Helianthus tuberosus</i> L. regardless of their presentation.

5. Other seeds and fruits, and products derived thereof

Number	Name	Description
5.1.1	Acorn	Whole fruits of the pendunculate oak <i>Quercus robur</i> L., the sessile oak <i>Quercus petraea</i> (Matt.) Liebl., the cork oak of <i>Quercus suber</i> L., or other species of oak.
5.1.2	Acorn, dehulled	Product obtained during dehulling of acorn.
5.2.1	Almond	Whole or broken fruit <i>Prunus dulcis</i> , with or without hulls.
5.2.2	Almond Hulls	Almond hulls obtained from dehusked almond seeds by physical separation from the kernels and ground.
5.2.3	Almond kernel expeller	Product of oil manufacture obtained by pressing of almond kernels
5.3.1	Anise seed	Seeds of <i>Pimpinella anisum</i> .
5.4.1	Apple pulp, dried; [Apple pomace, dried]	Product obtained from the production of juice of <i>Malus domestica</i> or cider production. It consists principally of internal pulp and outer skins that are dried. It may have been depectinised
5.4.2	Apple pulp, pressed; [Apple pomace, pressed]	Moist product obtained from the production of apple juice or cider production. It consists principally of internal pulp and outer skins that are pressed. It may have been depectinised
5.4.3	Apple molasses	Product obtained after producing pectin from apple pulp. It may have been depectinised
5.5.1	Sugar beet seed	Seeds of sugar beet
5.6.1	Buckwheat	Seeds of <i>Fagopyrum esculentum</i>
5.6.2	Buckwheat hulls and bran	Product obtained during the milling of buckwheat grains.
5.6.3	Buckwheat middlings	Product of flour manufacture, obtained from screened buckwheat. It consists principally of particles of endosperm, with fine fragments of the outer and some miscellaneous parts of the grain. It must contain no more than 10% crude fibre.
5.7.1	Red cabbage seed	Seeds of <i>Brassica oleracea</i> var. <i>capitata</i> f. <i>Rubra</i> .
5.8.1	Canary grass seed	Seeds of <i>Phalaris canariensis</i> .
5.9.1	Caraway seed	Seeds from <i>Carum carvi</i> L.
5.12.1	Broken chestnuts	Product of the production of chestnut flour, consisting mainly of particles of endosperm, with fine fragments of envelopes and a few remnants of chestnut (<i>Castanea</i> spp.).
5.13.1	Citrus pulp	Product obtained by pressing citrus fruits <i>Citrus</i> (L.) spp. or during the production of citrus juice. It may have been depectinised.
5.13.2	Citrus pulp, dried	Product obtained by pressing citrus fruits or during the production of citrus juice, which is subsequently dried. It may have been depectinised.
5.14.1	Red clover seed	Seeds of <i>Trifolium pratense</i> L.
5.14.2	White clover seed	Seeds of <i>Trifolium repens</i> L.
5.15.1	Coffee skins	Product obtained from dehusked seeds of the <i>Coffea</i> plant.
5.16.1	Cornflower seed	Seeds of <i>Centaurea cyanus</i> L.
5.17.1	Cucumber seed	Seeds of <i>Cucumis sativus</i> L.
5.18.1	Cypress seed	Seeds of <i>Cupressus</i> L.
5.19.1	Date fruit	Fruits of <i>Phoenix dactylifera</i> L. It may be dried.
5.19.2	Date seed	Whole seeds of the date plant.
5.20.1	Fennel seed	Seeds of <i>Foeniculum vulgare</i> Mill.
5.21.1	Fig fruit	Fruits of <i>Ficus carica</i> L. It may be dried.
5.22.1	Fruit kernels	Product consisting of the inner, edible seeds of a nut or fruit stone.
5.22.2	Fruit pulp	Product obtained during the production of fruit juice and fruit puree. It may have been depectinised.
5.22.3	Fruit pulp, dried	Product obtained during the production of fruit juice and fruit puree which is subsequently dried. It may have been depectinised.
5.23.1	Garden cress	Seeds from <i>Lepidium sativum</i> L.
5.24.1	Graminaceous seeds	Seeds from graminoids of the families <i>Poaceae</i> , <i>Cyperaceae</i> and <i>Juncaceae</i> .
5.25.1	Grape pips	Pips from <i>vitis</i> L. separated from grape pulp, from which the oil has not been removed.
5.25.2	Grape pips meal	Product obtained during the extraction of oil from grape pips.
5.25.3	Grape pulp [Grape marc]	Grape pulp dried rapidly after the extraction of alcohol from which as much as possible of the stalks and pips have been removed.
5.25.4	Grape pips soluble	Product obtained from grape pips after producing grape juice. It principally contains carbohydrates. It may be concentrated.
5.26.1	Hazelnut	Whole or broken fruit of <i>Corylus</i> (L.) spp., with or without hulls.
5.26.2	Hazelnut expeller	Product of oil manufacture obtained by pressing of hazelnut kernels

5.27.1	Pectin	Pectin is obtained by aqueous extraction (of natural strains) of appropriate plant material, usually citrus fruits or apples. No organic precipitant shall be used other than methanol, ethanol and propane-2-ol. May contain up to 1% methanol, ethanol and propane-2-ol singly or in combination, on an anhydrous basis. Pectin consists mainly of the partial methyl esters of polygalacturonic acid and their ammonium, sodium, potassium and calcium salts.
5.28.1	Perilla seed	Seeds of <i>Perilla frutescens</i> L. and its milling products.
5.29.1	Pine nut	Seeds from <i>Pinus</i> (L.) spp.
5.30.1	Pistachio	Fruit of <i>Pistacia vera</i> L.
5.31.1	Plantago seed	Seeds of <i>Plantago</i> (L.) spp.
5.32.1	Radish seed	Seeds of <i>Raphanus sativus</i> L.
5.33.1	Spinach seed	Seeds of <i>Spinacia oleracea</i> L.
5.34.1	Thistle seed	Seeds from <i>Carduus marianus</i> L.
5.35.1	Tomato pulp [tomato pomace]	Product obtained by pressing tomatoes <i>Solanum lycopersicum</i> L. during the production of tomato juice. It consists principally of tomato peel and seeds.
5.36.1	Yarrow seed	Seeds of <i>Achillea millefolium</i> L.
5.37.1	Apricot kernel expeller	Product of oil manufacture obtained by pressing of apricot kernels (<i>Prunus armeniaca</i> L.). It may contain hydrocyanic acid
5.38.1	Black cumin expeller	Product of oil manufacture obtained by pressing of black cumin seeds (<i>Bunium persicum</i> L.)
5.39.1	Borage seed expeller	Product of oil manufacture obtained by pressing of borage seeds (<i>Borago officinalis</i> L.)
5.40.1	Evening primrose expeller	Product of oil manufacture obtained by pressing of evening primrose seeds (<i>Oenothera</i> L.)
5.41.1	Pome grenade expeller	Product of oil manufacture obtained by pressing of pome grenade seeds (<i>Punica granatum</i> L.)
5.42.1	Walnut kernel expeller	Product of oil manufacture obtained by pressing of walnut kernels (<i>Juglans regia</i> L.)

6. Forages and roughage, and products derived thereof

Number	Name	Description
6.1.1	Beet leaves	Leaves of <i>Beta</i> spp.
6.2.1	Cereal plants	Whole plants of cereal species or parts thereof. It may be dried, fresh or ensiled.
6.3.1	Cereals straw	Straw of cereals.
6.3.2	Cereal straw, treated	Product obtained by an appropriate treatment of cereal straw.
6.4.1	Clover meal	Product obtained by drying and milling clover <i>Trifolium</i> spp. It may contain up to 20% lucerne (<i>Medicago sativa</i> L. and <i>Medicago</i> var. <i>Martyn</i>) or other forage crops dried and milled at the same time as the clover.
6.5.1	Forage meal; [Grass meal]; [Green meal]	Product obtained by drying and milling and in some cases compacting forage plants.
6.6.1	Grass, field dried, [Hay]	Species of any grass, field dried.
6.6.2	Grass, high temperature dried	Product obtained from grass (any variety) that has been artificially dehydrated (in any form).
6.6.3	Grass, herbs, legume plants, [green forage]	Fresh, ensiled or dried arable crops consisting of grass, legumes or herbs, commonly described as silage, haylage, hay or green forage.
6.7.1	Hemp flour	Flour ground from dried leaves from <i>Cannabis sativa</i> L.
6.7.2	Hemp fibre	Product obtained during the processing of hemp, green coloured, dried, fibrous.
6.8.1	Horse bean straw	Straw of horse bean.
6.9.1	Linseed straw	Straw of linseed (<i>Linum usitatissimum</i> L.).
6.10.1	Lucerne; [Alfalfa]	<i>Medicago sativa</i> L. and <i>Medicago</i> var. <i>Martyn</i> plants or parts thereof.
6.10.2	Lucerne field dried; [Alfalfa field dried]	Lucerne, field dried.
6.10.3	Lucerne, high temperature dried; [Alfalfa, high temperature dried]	Lucerne artificially dehydrated, in any form.
6.10.4	Lucerne, extruded; [Alfalfa, extruded]	Alfalfa pellets that have been extruded.
6.10.5	Lucerne meal; [Alfalfa meal]	Product obtained by drying and milling Lucerne. It may contain up to 20% clover or other forage crop dried and milled at the same time as the lucerne.
6.10.6	Lucerne pomace; [Alfalfa pomace]	Dried product obtained by pressing of the juice from lucerne.
6.10.7	Lucerne protein concentrate; [Alfalfa protein concentrate]	Product obtained by artificially drying fractions of lucerne press juice, which have been separated by centrifugation and heat treated to precipitate the proteins.
6.10.8	Lucerne solubles	Product obtained after the extraction of proteins from lucerne juice, it may be dried
6.11.1	Maize silage	Ensiled plants or parts thereof of <i>Zea mays</i> L. ssp. <i>mays</i> .
6.12.1	Pea Straw	Straw of <i>Pisum</i> spp.

7. Other plants, algae and products derived thereof

Number	Name	Description
7.1.1	Algae	Algae, live or processed, including fresh, chilled or frozen algae. May contain up to 0.1% of antifoaming agents.
7.1.2	Dried algae	Product obtained by drying algae. This product may have been washed to reduce the iodine content. May contain up to 0.1% of antifoaming agents.
7.1.3	Algae meal	Product of algae oil manufacture, obtained by extraction of algae. May contain up to 0.1% of antifoaming agents.
7.1.4	Algal oil	Product of the oil manufacture from algae obtained by extraction. May contain up to 0.1% of antifoaming agents.
7.1.5	Algae extract; [Algae fraction]	Watery or alcoholic extract of algae that principally contains carbohydrates. May contain up to 0.1% of antifoaming agents.
7.2.6	Seaweed meal	Product obtained by drying and crushing macro-algae, in particular brown seaweed. This product may have been washed to reduce the iodine content. May contain up to 0.1% of antifoaming agents.
7.3.1	Barks	Cleaned and dried barks of trees or bushes.
7.4.1	Blossoms, dried	All parts of dried blossoms of consumable plants and their fractions.
7.5.1	Broccoli, dried	Product obtained by drying the plant <i>Brassica oleracea</i> L. after washing, size reduction (cutting, flaking, etc.) and water content removal.
7.6.1	(Sugar) cane molasses	Syrupy product obtained during the manufacture or refining of sugar from <i>Saccharum</i> L. May contain up to 0.5% antifoaming agents. May contain up to 0.5% antiscaling agents. May contain up to 3.5% sulphate. May contain up to 0.25% sulphite.
7.6.2	(Sugar) cane Molasses, partially desugared	Product obtained after further extraction using water of sucrose from sugar cane molasses.
7.6.3	(Cane) sugar [sucrose]	Sugar extracted from sugar canes using water.
7.6.4	Cane bagasse	Product obtained during extraction using water of sugar from sugar canes. It consists mainly of fibres.
7.7.1	Leaves, dried	Dried leaves of consumable plants and their fractions.
7.8.1	Lignocellulose	Product obtained by means of mechanical processing of raw natural dried wood and which predominantly consists of lignocellulose.
7.9.1	Liquorice root	Root of <i>Glycyrrhiza</i> L.
7.10.1	Mint	Product obtained from drying aerial parts of the plants <i>Mentha apicata</i> , <i>Mentha piperita</i> or <i>Mentha viridis</i> (L.), regardless of their presentation.
7.11.1	Spinach, dried	Product obtained from drying the plant <i>Spinacia oleracea</i> L., regardless of its presentation.
7.12.1	Mojave yucca	Pulverized <i>Yucca schidigera</i> Roezl.
7.13.1	Vegetal carbon; [charcoal]	Product obtained by carbonisation of organic vegetal material.
7.14.1	Wood	Chemically untreated mature wood or wood fibres.

11. Minerals

Number	Name	Description
11.1.1	Calcium carbonate; [Limestone]	Product obtained by grinding sources of calcium carbonate (CaCO ₃), such as limestone or by precipitation from acid solution. May contain up to 0.25% propylene glycol. May contain up to 0.1% grinding aids.
11.1.2	Calcareous marine shells	Product of natural origin, obtained from marine shells, ground or granulated, such as oyster shells or seashells.
11.1.3	Calcium and magnesium carbonate	Natural mixture of calcium carbonate (CaCO ₃) and magnesium carbonate (MgCO ₃). May contain up to 0.1% grinding aids.
11.1.4	Maerl	Product of natural origin obtained from calcareous marine algae, ground or granulated.
11.1.5	Lithothamn	Product of natural origin obtained from calcareous marine algae (<i>Phymatolithon calcareum</i> (Pall.)), ground or granulated.
11.1.6	Calcium chloride	Calcium chloride (CaCl ₂). May contain up to 0.2% barium sulphate
11.1.7	Calcium hydroxide	Calcium hydroxide (Ca(OH) ₂). May contain up to 0.1% grinding aids.
11.1.8	Calcium sulphate anhydrous	Calcium sulphate anhydrous (CaSO ₄) obtained by grinding calcium sulphate anhydrous or dehydration of calcium sulphate dihydrate.
11.1.9	Calcium sulphate hemihydrate	Calcium sulphate hemihydrate (CaSO ₄ x ½ H ₂ O) obtained by partially dehydrating calcium sulphate dihydrate.
11.1.10	Calcium sulphate dihydrate	Calcium sulphate dihydrate (CaSO ₄ x 2H ₂ O) obtained by grinding calcium sulphate dihydrate or hydration of calcium sulphate hemihydrate.
11.1.11	Calcium salts of organic acids	Calcium salts of edible organic acids with at least 4 carbon atoms.
11.1.12	Calcium oxide	Calcium oxide (CaO) obtained from calcination of naturally occurring limestone. May contain up to 0.1% grinding aids.
11.1.13	Calcium gluconate	Calcium salt of gluconic acid generally expressed as Ca(C ₆ H ₁₁ O ₇) ₂ and its

		hydrated forms.
11.1.15	Calcium Sulphate / Carbonate	Product obtained during the manufacturing of sodium carbonate.
11.1.16	Calcium pidolate	L-calcium pidolate ($C_5H_6CaNO_3$). May contain up to 1.5% glutamic acid and related substances.
11.1.17	Calcium carbonate-magnesium oxide	Product obtained by heating of natural calcium and magnesium containing substances like dolomite. May contain up to 0.1% grinding aids.
11.2.1	Magnesium oxide	Calcined magnesium oxide (MgO) not less than 70% MgO .
11.2.2	Magnesium sulphate heptahydrate	Magnesium sulphate ($MgSO_4 \times 7 H_2O$).
11.2.3	Magnesium sulphate monohydrate	Magnesium sulphate ($MgSO_4 \times H_2O$).
11.2.4	Magnesium sulphate anhydrous	Anhydrous magnesium sulphate ($MgSO_4$).
11.2.5	Magnesium propionate	Magnesium propionate ($C_6H_{10}MgO_4$).
11.2.6	Magnesium chloride	Magnesium chloride ($MgCl_2$) or solution obtained by natural concentration of sea water after deposit of sodium chloride.
11.2.7	Magnesium carbonate	Natural magnesium carbonate ($MgCO_3$).
11.2.8	Magnesium hydroxide	Magnesium hydroxide ($Mg(OH)_2$).
11.2.9	Magnesium potassium sulphate	Magnesium potassium sulphate.
11.2.10	Magnesium salts of organic acids	Magnesium salts of edible organic acids with at least 4 carbon atoms.
11.3.1	Dicalcium phosphate; [Calcium hydrogen orthophosphate]	Calcium mono-hydrogen phosphate obtained from bones or inorganic sources ($CaHPO_4 \times H_2O$) $Ca/P > 1.2$ May contain up to 3% chloride expressed as NaCl
11.3.2	Mono-dicalcium phosphate	Product obtained chemically and composed of dicalcium phosphate and mono-calcium phosphate ($CaHPO_4, Ca(H_2PO_4)_2 \times H_2O$) $0.8 < Ca/P < 1.3$
11.3.3	Mono-calcium phosphate; [Calcium tetrahydrogen diorthophosphate]	Calcium-bis dihydrogenphosphate ($Ca(H_2PO_4)_2 \times H_2O$) $Ca/P < 0.9$
11.3.4	Tri-calcium phosphate; [Tricalcium orthophosphate]	Tri-calcium phosphate from bones or inorganic sources ($Ca_3(PO_4)_2 \times H_2O$) $Ca/P > 1.3$
11.3.5	Calcium-magnesium phosphate	Calcium-magnesium phosphate.
11.3.6	Defluorinated phosphate	Natural phosphate, calcined and further heat treated than for the removal of impurities necessary.
11.3.7	Di-calcium pyrophosphate; [Dicalcium diphosphate]	Dicalcium pyrophosphate.
11.3.8	Magnesium phosphate	Product consisting of monobasic and/or di-basic and/or tri-basic magnesium phosphate.
11.3.9	Sodium-calcium-magnesium phosphate	Product consisting of sodium-calcium-magnesium phosphate.
11.3.10	Mono-sodium phosphate; [Sodium dihydrogen orthophosphate]	Mono-sodium phosphate ($NaH_2PO_4 \times H_2O$)
11.3.11	Di-sodium phosphate; [Disodium hydrogen orthophosphate]	Disodium phosphate ($Na_2HPO_4 \times H_2O$)
11.3.12	Tri-sodium Phosphate; [Trisodium orthophosphate]	Tri-sodium phosphate (Na_3PO_4)
11.3.13	Sodium pyrophosphate; [Tetrasodium diphosphate]	Sodium pyrophosphate ($Na_4P_2O_7$).
11.3.14	Mono-potassium phosphate; [Potassium dihydrogen orthophosphate]	Mono-potassium phosphate ($KH_2PO_4 \times H_2O$)
11.3.15	Di-potassium phosphate; [Di-potassium hydrogen orthophosphate]	viii. Di-potassium phosphate ($K_2HPO_4 \times H_2O$)
11.3.16	Calcium sodium phosphate	Calcium sodium phosphate ($CaNaPO_4$).
11.3.17	Mono-ammonium phosphate; [Ammonium dihydrogen orthophosphate]	Mono-ammonium phosphate ($NH_4H_2PO_4$)
11.3.18	Di-ammonium phosphate; [Diammonium hydrogen orthophosphate]	Di-ammonium phosphate ($(NH_4)_2HPO_4$)
11.3.19	Sodium tri-polyphosphate; [Penta sodium triphosphate]	Sodium tri-polyphosphate ($Na_5P_3O_9$).
11.3.20	Sodium magnesium phosphate	Sodium-magnesium phosphate ($MgNaPO_4$).
11.3.21	Magnesium hypophosphite	Magnesium hypophosphite ($Mg(H_2PO_2)_2 \times 6H_2O$)
11.3.22	Degelatinised bone meal	Degelatinised, sterilised and ground bones from which the fat has been removed.
11.3.23	Bone ash	Mineral residues from the incineration, combustion or gasification of animal by-

		products.
11.3.24	Calcium polyphosphate	Heterogeneous mixtures of calcium salts of condensed polyphosphoric acids of general formula $H(n+2)PnO(3n+1)$ where 'n' is not less than 2
11.3.25	Calcium dihydrogen diphosphate	Mono-calcium dihydrogen pyrophosphate ($CaH_2P_2O_7$).
11.3.26	Magnesium acid pyrophosphate	Magnesium acid pyrophosphate ($MgH_2P_2O_7$). Produced from purified phosphoric acid and purified magnesium hydroxide or magnesium oxide by evaporation of water and condensation of the orthophosphate to diphosphate.
11.3.27	Disodium dihydrogen diphosphate	Disodium dihydrogen diphosphate ($Na_2H_2P_2O_7$).
11.3.28	Tri-sodium diphosphate	Trisodium monohydrogen diphosphate (anhydrous: $Na_3HP_2O_7$; monohydrate: $Na_3HP_2O_7 \times H_2O$)
11.3.29	Sodium polyphosphate; [Sodium hexametaphosphate]	Heterogeneous mixtures of sodium salts of linear condensed polyphosphoric acids of general formula $H(n+2)PnO(3n+1)$ where 'n' is not less than 2
11.3.30	Tri-potassium phosphate	Tripotassium monophosphate (anhydrous: K_3PO_4 ; hydrated: $K_3PO_4 \times n H_2O$ (n=1 or 3)).
11.3.31	Tetrapotassium di-phosphate	Tetrapotassium pyrophosphate ($K_4P_2O_7$)
11.3.32	Pentapotassium tri-phosphate	Pentapotassium tri-polyphosphate ($K_5P_3O_{10}$)
11.3.33	Potassium polyphosphate	Heterogeneous mixtures of potassium salts of linear condensed polyphosphoric acids of general formula $H(n+2)PnO(3n+1)$ where 'n' is not less than 2
11.3.34	Calcium sodium polyphosphate	Calcium sodium polyphosphate.
11.4.1	Sodium chloride	Sodium chloride ($NaCl$) or product obtained by evaporative crystallization from brine (vacuum salt) or evaporation of seawater (marine salt) or grinding rock salt.
11.4.2	Sodium bicarbonate [Sodium hydrogencarbonate]	Sodium bicarbonate ($NaHCO_3$)
11.4.3	Sodium/ammonium (bi)carbonate [Sodium/ammonium (hydrogen)carbonate]	Product obtained during the production of sodium carbonate and sodium bicarbonate, with traces of ammonium bicarbonate (ammonium bicarbonate max. 5%).
11.4.4	Sodium carbonate	Sodium carbonate (Na_2CO_3)
11.4.5	Sodium sesquicarbonate [Trisodium hydrogencarbonate]	Sodium sesquicarbonate ($Na_3H(CO_3)_2$)
11.4.6	Sodium sulphate	Sodium sulphate (Na_2SO_4). May contain up to 0.3% methionine
11.4.7	Sodium salts of organic acids	Sodium salts of edible organic acids with at least 4 carbon atoms.
11.5.1	Potassium chloride	Potassium chloride (KCl) or product obtained by grinding natural sources of potassium chloride.
11.5.2	Potassium sulphate	Potassium sulphate (K_2SO_4)
11.5.3	Potassium carbonate	Potassium carbonate (K_2CO_3).
11.5.4	Potassium bicarbonate [Potassium hydrogen carbonate]	Potassium bicarbonate ($KHCO_3$).
11.5.5	Potassium salts of organic acids	Potassium salts of edible organic acids with at least 4 carbon atoms.
11.6.1	Flower of sulphur	Powder obtained from natural deposits of the mineral. Also, product obtained from oil refinery production as practised by sulphur manufacturers.
11.7.1	Attapulgite	Natural magnesium-aluminium-silicon mineral.
11.7.2	Quartz	Naturally occurring mineral obtained by grinding sources of quartz. May contain up to 0.1% grinding aids.
11.7.3	Cristobalite	Silicon dioxide (SiO_2) obtained from the re-crystallisation of quartz. May contain up to 0.1% grinding aids.
11.8.1	Ammonium sulphate	Ammonium sulphate ($(NH_4)_2SO_4$) obtained by chemical synthesis.
11.8.2	Ammonium sulphate solution	Ammonium sulphate in aqueous solution, containing not less than 35% Ammonium sulphate
11.8.3	Ammonium salts of organic acids	Ammonium salts of edible organic acids with at least 4 carbon atoms.
11.8.4	Ammonium lactate	Ammonium lactate ($CH_3CHOHCOONH_4$). Includes the Ammonium lactate produced by fermentation with <i>Lactobacillus delbrueckii ssp. Bulgaricus</i> , <i>Lactococcus lactis ssp.</i> , <i>Leuconostoc mesenteroides</i> , <i>Streptococcus thermophilus</i> , <i>Lactobacillus spp.</i> , or <i>Bifidobacterium spp.</i> , containing not less than 44% Nitrogen expressed as crude protein. May contain up to 0.8% phosphorus, 0.9% potassium, 0.7% magnesium, 0.3% sodium, 0.3% sulphates 0.1% chlorides, 5% sugars and 0.1% silicone antifoam.
11.8.5	Ammonium acetate	Ammonium acetate (CH_3COONH_4) in aqueous solution, containing not less than 55% Ammonium acetate)

These materials may be defined further on the label by processes they have undergone, such as: dried, flaked, extruded, flour, toasted, pelleted etc. Thus examples of feed materials as listed on a label would be: dried peas, extruded wheat and flaked maize.

If a feed material does not correspond exactly with the definition given above, an alternative accurate description of the feed material should be used. Several other feed materials not in the EU catalogue are also legitimately used for rabbit foods, provided they are safe and accurately described on the label. A few examples of additional feed materials are given in Table 2.

Some examples of additional feed materials for rabbit food not in the EU Catalogue, but which are included in the Register of Feed Materials (<http://www.feedmaterialsregister.eu/>)

Name	Description
Hay or Meadow Hay	Dried grass or sedge cut from meadows and dried.
Timothy hay	Dried grass of the species <i>Phleum pratense</i> .
Apple	Fruit of the tree <i>Malus domestica</i> .
Carob meal, extruded	Seeds and pods of the carob tree, <i>Ceratonia siliqua</i> , dried, ground and extruded.
Parsley, dried or oil	Parsley, <i>Petroselinum</i> sp. leaves; dried or the extracted oil
Herbs	Several herbs may be used in rabbit foods including Anise, <i>Pimpinella anisum</i> ; Oregano, <i>Oregano vulgare</i> ; Rosemary, <i>Rosmarinus officinalis</i> and others
Cellulose fibre	Pulp made from wood is dissolved and purified and dried to make cellulose. Cellulose with varying lengths of fibre is available and can be a source of fibre in the diet.
Spinach	Leaves of the species <i>Spinacia oleracea</i> ,
Dandelion	Leaves, and sometimes flower heads, of the species <i>Taraxacum officinale</i> , or the genus <i>Taraxacum</i> . Often included in a food as dried dandelion.

Bibliography

Balls, M., van Zeller, A.M., Halder, M.E. (1999) 'Information Resources on the Care and Welfare of Rabbits' *Elsevier Science B.V.*, 1269-1277.

Baylos, M., Menoyo, D., Chamorro, S., Sainz, A., Nicodemus, N., de Blas, J.C. and Carabano, R. (2008) 'Effect of dietary level and source of glutamine on intestinal health in the postweaning period' *Nutrition and Digestive Physiology*, 529 – 534.

Berthelsen, H. and Hansen, L.T. (1999) 'The effect of hay on the behaviour of caged rabbits (*Oryctolagus cuniculi*)' *Animal Welfare* 8 (2), 149-157.

Bilko, A., Altbacker, V., Hudson, R. (1994) 'Transmission of food preference in the rabbit: the means of information transfer' *Physiology of Behaviour* 56 (5), 907-912.

Blaxter, K.L. (1989) 'Energy Metabolism in Animals and Man' *Cambridge University Press*.

Bouyssou, T., Candau, M. and Ruckebush, Y. (1988) 'Réponses motrices du côlon aux constituants pariétaux et à la finesse de mouture des aliments chez le lapin. (Colonic motility pattern according to the source of fibre and to the grinding level of the diet, in the rabbit.)' *Reproduction Nutrition Développement* 28, 181–182.

Brooks, D.L., Huls, W., Leamon, C., Thomson, J., Parker, J. and Twomey, S. (1993) 'Cage enrichment for female New Zealand White rabbits' *Lab Animal*, 22: 30-35.

Chamorro, S., de Blas, C., Grant, G., Badiola, I., Menoyo, D. and Carabaño, R. (2010) 'Effect of dietary supplementation with glutamine and a combination of glutamine-arginine on intestinal health in twenty-five-day-old weaned rabbits' *Journal of Animal Science* 88, 170-180.

Cheeke, P.R. (1994) 'Nutrition and Nutritional Diseases' In: *The Biology of the Laboratory Rabbit* 2nd ed. *Academic Press, Orlando*, 321-333.

De Blas, C., Wiseman, J. (2010) 'Nutrition of the Rabbit' *CABI Publishing (2nd Revised edition) Wallingford, Oxford*.

Gidenne, T. and Garcia, J. (2006) 'Nutritional strategies improving the digestive health of the weaned rabbit' *Recent Advances in Rabbit Sciences' Ilvo, Merelbeke, Belgium*, 211-227.

Gunn, D. (1994) 'Evaluation on welfare in the husbandry of laboratory rabbits (PhD Thesis)' *University of Birmingham*.

Hall, M.B., et al (1997) A simple method for estimation of Neutral detergent soluble fibre. *J food Food Sci Agric* 74 441-449 Harcourt-Brown, F.M. (1996) 'Calcium deficiency, diet and dental disease in pet rabbits' *Veterinary Record* 139, 567-571.

Harcourt-Brown, F.M. (2002) 'Diet and Husbandry. In: Textbook of Rabbit Medicine' *Butterworth Heinemann, Oxford*, 19-51.

INRA 2004, Tables of composition and nutritional value of feed materials (Pigs, poultry, cattle, sheep, goats, rabbits, horses and fish), edited by: D. Sauvant, J.-M. Perez and G. Tran, Wageningen Academic Publishers, 2004

Jackson, G. (1991) 'Intestinal stasis and rupture in rabbits' *Veterinary Record* 129, 287-289.

Kamphues, J., Coenen, M. and Kienzle, E. (2009) 'Empfehlungen für den Energie- und

Nährstoffgehalt in Alleinfuttermitteln für Versuchstiere' *Supplemente zu Vorlesungen und Übungen in der Tierernährung*. 11. Aufl. M.& H. Schaper, S. 316

Lebas, F., (2000) 'Vitamins in rabbit nutrition : Literature review and recommendations' *World Rabbit Science* 2000, 8 (4), 185 – 192.

Lidfors, L. (1997) 'Behavioural effects of environmental enrichment for individually caged rabbits' *Applied Animal Behaviour Science* 52 (1), 157-169

Mason, G. and Rushen, J. (2006) 'Stereotypic Animal Behaviour: Fundamentals and Applications to Welfare' 2nd Edition' *CABI, Wallingford, Oxford*.

McDonald P, Edwards RA, Greenhalgh JFD, et al., (1995) 'Animal Nutrition' 7th edition. Longman GroupPearson Education Ltd, Harlow, UK.

Morton, D.B. (2002) 'Behaviour of rabbits and rodents. In: The Ethology of Domestic Animals: an Introductory Text' *CABI Publishing, Wallingford, Oxon*, 193-209.

Morton, D.B., Jennings, M., Batchelor, G.R., Bell, D., Birke, L., Davies, K., Eveleigh, J.R., Gunn, D., Heath, M., Howard, B., Koder, P., Phillips, J., Poole, T., Sainsbury, A.W., Sales, G.D., Smith, D.J.A., Stauffacher, M. and Turner, R.J. (1993) 'Refinements in rabbit husbandry' *Second report of the BVA/WF/FRAME/RSPCA/UFPA joint working group on refinement. Laboratory Animals* 27, 301-329.

Odberg, F. O. (1978) 'Abnormal behaviours: stereotypes' Paper presented at the *Proceedings of the 1st World Congress on Ethology Applied to Zootechnics*, Madrid.

Oswalt, M. and Kemp, S.F. (2007) 'Anaphylaxis: office management and prevention' *Immunology And Allergy Clinics of North America*, 27 (2) 177-191.

Pairet, M., Bouyssou, T. and Ruckebusch, Y. (1986) 'Colonic formation of soft faeces in rabbits: a role for endogenous prostaglandins' *American Journal Physiology* 250 (Gastrointestinal. Liver Physiology 13), G302-G308.

Reeds, P. and Beckett, P. (1996), 'Protein and amino acids' *Present Knowledge in Nutrition*. 7th ed. Washington, DC: International Life Sciences Institute; 67-86.

Stauffacher, M., (1992) 'Group housing and enrichment cages for breeding, fattening and laboratory rabbits' *Animal Welfare* 1: 105-125.

Stauffacher, M. (2000) 'Refinement in rabbit housing and husbandry' Progress in the Reduction, Refinement and Replacement of Animal Experimentation: *Proceedings of the 3rd World Congress on Alternatives and Animal Use in the Life Sciences*.

Tang, A.W. (2003) 'A practical guide to anaphylaxis' *Am Fam Physician* 68 (7), 1325-1332.

Tobin, G. (1996) Small Pets, Food types Nutrient Requirements and nutritional disorders in Manual of companion animal nutrition and feeding. Eds: Kelly, N and Wills, J.

Villamide, M.J., Carabaño, R., Maertens, L., Pascual, J., Gidenne, T., Falcao-E-Cunha, L. and Xiccato, G. (2009) 'Prediction of the nutritional value of European compound feeds for rabbits by chemical components and *in vitro* analysis' *Animal Feed Science and Technology* 150, 283-295.

Wallace, S., Sanford, J., Smith, M. W. and Spencer, K. V. (1990) 'The assessment and control of the severity of scientific procedures on laboratory animals - Report of the LASA Working Party

(Assessment and control of severity)' *Laboratory Animals* 24, 97-130.

Wang, J., Sampson, H.A. (2007) 'Food Anaphylaxis' *Clinical and Experimental Allergy*, 37 (5), 651-660.

* * * *